

09/784,554

**PCT**WORLD INTELLECTUAL PROPERTY ORGANIZATION  
International Bureau

## INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

<b>(51) International Patent Classification <sup>6</sup> :</b> <b>C12N 9/42 // 9/42, C12R 1:07</b>	<b>A1</b>	<b>(11) International Publication Number:</b> <b>WO 99/02663</b> <b>(43) International Publication Date:</b> 21 January 1999 (21.01.99)
<b>(21) International Application Number:</b> PCT/DK98/00290 <b>(22) International Filing Date:</b> 1 July 1998 (01.07.98)  <b>(30) Priority Data:</b> 0822/97                      7 July 1997 (07.07.97)                      DK 1213/97                      24 October 1997 (24.10.97)                      DK  <b>(71) Applicant:</b> NOVO NORDISK A/S [DK/DK]; Novo Allé, DK-2880 Bagsvaerd (DK).  <b>(72) Inventors:</b> SCHÜLEIN, Martin; Wiedeweltsgade 51, DK-2100 Copenhagen Ø (DK). OUTTRUP, Helle; Syvendehusvej 46, DK-2750 Ballerup (DK). JØRGENSEN, Per, Linå; Dr. Tvaergade 37, i., DK-1302 Copenhagen K (DK). BJØRNVAD, Mads, Eskelund; Dr. Abildgaards Allé 8, 3th, DK-1955 Frederiksberg (DK).  <b>(74) Common Representative:</b> NOVO NORDISK A/S; Corporate Patents, Novo Allé, DK-2880 Bagsvaerd (DK).		<b>(81) Designated States:</b> AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, CA, CH, CN, CU, CZ, DE, DK, EE, ES, FI, GB, GE, GH, GM, GW, HR, HU, ID, IL, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MD, MG, MK, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, UA, UG, UZ, VN, YU, ZW, ARIPO patent (GH, GM, KE, LS, MW, SD, SZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, ML, MR, NE, SN, TD, TG).  <b>Published</b> <i>With international search report.</i>
<b>(54) Title:</b> ALKALINE XYLOGLUCANASE  <b>(57) Abstract</b>  <p>A xyloglucanase having a relative xyloglucanase activity of at least 50% at pH7 and either no or an insignificant cellulolytic activity is obtainable e.g. from a strain of <i>Bacillus</i>. A xyloglucanase comprising an amino acid sequence as shown in positions 30-261 of SEQ ID NO:2 or homologues may be derived from e.g. <i>Bacillus licheniformis</i>, ATCC 14580, and may be encoded by polynucleotide molecules comprising a nucleotide sequence as shown in SEQ ID NO:1 from nucleotide 88 to nucleotide 783; and a xyloglucanase comprising an amino acid sequence as shown in positions 1-537 of SEQ ID NO:4 or homologues may be derived from e.g. <i>B. agaradhaerens</i>, NCIMB 40482, and may be encoded by polynucleotide molecules comprising a nucleotide sequence as shown in SEQ ID NO:3 from nucleotide 1 to nucleotide 1611. The xyloglucanases are useful e.g. in cleaning compositions and for treatment of cellulosic fibres.</p>		

**FOR THE PURPOSES OF INFORMATION ONLY**

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AL	Albania	ES	Spain	LS	Lesotho	SI	Slovenia
AM	Armenia	FI	Finland	LT	Lithuania	SK	Slovakia
AT	Austria	FR	France	LU	Luxembourg	SN	Senegal
AU	Australia	GA	Gabon	LV	Latvia	SZ	Swaziland
AZ	Azerbaijan	GB	United Kingdom	MC	Monaco	TD	Chad
BA	Bosnia and Herzegovina	GE	Georgia	MD	Republic of Moldova	TG	Togo
BB	Barbados	GH	Ghana	MG	Madagascar	TJ	Tajikistan
BE	Belgium	GN	Guinea	MK	The former Yugoslav Republic of Macedonia	TM	Turkmenistan
BF	Burkina Faso	GR	Greece	ML	Mali	TR	Turkey
BG	Bulgaria	HU	Hungary	MN	Mongolia	TT	Trinidad and Tobago
BJ	Benin	IE	Ireland	MR	Mauritania	UA	Ukraine
BR	Brazil	IL	Israel	MW	Malawi	UG	Uganda
BY	Belarus	IS	Iceland	MX	Mexico	US	United States of America
CA	Canada	IT	Italy	NE	Niger	UZ	Uzbekistan
CF	Central African Republic	JP	Japan	NL	Netherlands	VN	Viet Nam
CG	Congo	KE	Kenya	NO	Norway	YU	Yugoslavia
CH	Switzerland	KG	Kyrgyzstan	NZ	New Zealand	ZW	Zimbabwe
CI	Côte d'Ivoire	KP	Democratic People's Republic of Korea	PL	Poland		
CM	Cameroon	KR	Republic of Korea	PT	Portugal		
CN	China	KZ	Kazakhstan	RO	Romania		
CU	Cuba	LC	Saint Lucia	RU	Russian Federation		
CZ	Czech Republic	LI	Liechtenstein	SD	Sudan		
DE	Germany	LK	Sri Lanka	SE	Sweden		
DK	Denmark	LR	Liberia	SG	Singapore		
EE	Estonia						

## ALKALINE XYLOGLUCANASE

The present invention relates to alkaline xyloglucanases, i.e. enzymes exhibiting xyloglucanase activity as their major enzymatic activity in the neutral and alkaline pH ranges; to a method of producing such enzymes; and to methods for using such enzymes in the textile, detergent and cellulose fiber processing industries.

## 10 BACKGROUND OF THE INVENTION

Xyloglucan is a major structural polysaccharide in the primary (growing) cell wall of plants. Structurally, xyloglucans consists of a cellulose-like beta-1,4-linked glucose backbone which is frequently substituted with various side chains. The xyloglucans of most dicotyledonous plants, some monocotyledons and gymnosperms are highly branched polysaccharides in which approx. 75% of the glucose residues in the backbone bear a glycosyl side chain at O-6. The glycosyl residue that is directly attached to the branched glucose residue is invariably alpha-D-xylose. Up to 50% of the side chains in the xyloglucans contain more than one residue due to the presence of beta-D-galactose or alpha-L-fucose-(1-2)-beta-D-galactose moieties at O-2 of the xylose residues (C. Ohsumi and T. Hayashi (1994) Plant and Cell Physiology 35:963-967; G. J. McDougall and S. C. Fry (1994) Journal of Plant Physiology 143:591-595; J. L. Acebes et al. (1993) Phytochemistry 33:1343-1345). On acid hydrolysis, the xyloglucan extracted from cotton fibers yielded glucose, xylose, galactose and fucose in the ratio of 50:29:12:7 (Hayashi et al., 1988).

30 Xyloglucans produced by solanaceous plants are unusual in that typical only 40% of the beta-1,4-linked glucose residues bear a glycosyl side chain at O-6. Furthermore, up to 60% of the xylose residues are substituted at O-2 with alpha-L-arabinose residues and some solanaceous plants, such as potato, also have xyloglucans with beta-D-galactose substituents at O-2 of some of the xylose residues (York et al (1996)).

Xyloglucan is believed to function in the primary wall of plants by crosslinking cellulose-microfibrils, forming a cellulose-xyloglucan network. This network is considered

necessary for the structural integrity of primary cell-walls (Carpita et al., 1993). Another important function of xyloglucan is to act as a repository for xyloglucan subunit oligo-saccharides that are physiologically active regulators of plant cell growth. Xyloglucan subunits may also modulate the action of a xyloglucan endotransglycosylase (XET), a cell-wall associated enzyme that has been hypothesized to play a role in the elongation of plant cell walls. Therefore xyloglucan might play an important role in wall loosening and consequently cell expansion (Fry et al., 1992).

The seeds of many dicotyledonous species contain xyloglucan as the major polysaccharide storage reserve. This type of xyloglucan, which is localized in massive thickenings on the inside of the seed cotyledon cell wall, is composed mainly of glucose, xylose and galactose (Rose et al., 1996).

Seeds of the tamarind tree *Tamarindus indica* became a commercial source of gum in 1943 when the gum was found useful as a paper and textile size. Sizing of jute and cotton with tamarind xyloglucan has been extensively practiced in Asia owing to the low cost of the gum and to its excellent properties. Food applications of tamarind xyloglucan include use in confections, jams and jellies and as a stabilizer in ice cream and mayonnaise (Whistler et al., 1993).

Xyloglucanase activity is not included in the classification of enzymes provided by the Enzyme Nomenclature (1992). Hitherto, this enzymatic activity has simply been classified as glucanase activity and has often been believed to be identical to cellulolytic activity (EC 3.2.1.4), i.e. activity against  $\beta$ -1,4-glycosidic linkages in cellulose or cellulose derivative substrates, or at least to be a side activity in enzymes having cellulolytic activity. However, a true xyloglucanase is a true xyloglucan specific enzyme capable of catalyzing the solubilisation of xyloglucan to xyloglucan oligosaccharides but which does not exhibit substantial cellulolytic activity, e.g. activity against the conventionally used cellulose-like substrates CMC (carboxymethylcellulose), HE cellulose and Avicel (microcrystalline cellulose). A xyloglucanase cleaves the beta-1,4-glycosidic linkages in the backbone of xyloglucan.

Xyloglucanase activity is described by Vincken et al. (1997) who characterizes three different endoglucanases from *Trichoderma viride* (similar to *T. reesei*) which all have high activity against cellulose or CMC and show that the EndoI (which is indeed belonging to family 5 of glycosyl hydrolases, see Henrissat, B. et al. (1991, 1993)) has essentially no (i.e. very little) activity against xyloglucan, and that EndoV (belonging to the family 7 of glycosyl hydrolases) and EndoIV (belonging to the family 12 of glycosyl hydrolases) both have activity against xyloglucan and CMC, respectively, of the same order of magnitude.

International patent publication WO 97/13862 describes two cellulases from *Aspergillus niger* strain N400 as described in EP-A 0 463 706. The sequence of lac12 can be ascribed to family 12, and the sequence of lac64 is determined as belonging to family 5 by comparison with known homologous cellulases in the EMBL data base. Both enzyme have cellulase and  $\beta$ -glucanase activity. They have the highest activity against barley  $\beta$ -glucan, and they both show good CMC activity and some xyloglucanase activity. Both enzyme has pH optima at 3.5 on CMC and 5.5 on barley  $\beta$ -glucan.

International Patent Publication WO 94/14953 discloses a xyloglucanase (EG II) cloned from the fungus *Aspergillus aculeatus* and expressed in the fungus *Aspergillus oryzae* which has high xyloglucanase activity and very little cellulase activity. This EG II enzyme which shows xyloglucanase activity in the pH range 2.5-6 and optimum activity at pH 3-4 also belongs to family 12 of glycosyl hydrolases.

In summary, up till now xyloglucanase activity has only been found in fungal enzymes belonging to the families 7 and 12 of glycosyl hydrolases and exhibiting this activity in the acidic to near neutral pH range.

However, many important processes, either industrial or using industrially produced agents, are operating at an alkaline pH. Thus, it is an object of the present invention to provide a true xyloglucanase enzyme with a high xyloglucanase activity at an alkaline pH and essentially no activity on cellulose or cellulose derivatives.

## SUMMARY OF THE INVENTION

The inventors have now found enzymes having substantial xyloglucanase activity in the alkaline range, such enzymes having either no or an insignificant cellulolytic activity.

5 Accordingly, the present invention relates to an enzyme preparation comprising a xyloglucanase having a relative xyloglucanase activity of at least 50% at pH 7 or a pH above 7, and preferably a minor or no activity on cellulose or cellulose derivative substrates, e.g. having a ratio of maximum  
10 xyloglucanase activity to maximum activity on CMC or Avicel of at least 2:1.

The inventors have also succeeded in cloning and expressing a xyloglucanase, ie the invention relates in further aspects to a xyloglucanase which is (a) a polypeptide produced by *Bacillus*  
15 *agaradhaerens*, N CIMB 40482, or (b) a polypeptide comprising an amino acid sequence as shown in positions 1-537 SEQ ID NO:4, or (c) an analogue of the polypeptide defined in (a) or (b) which is at least 70% homologous with said polypeptide, or is derived from said polypeptide by substitution, deletion or addition of  
20 one or several amino acids, or is immunologically reactive with a polyclonal antibody raised against said polypeptide in purified form; and to a xyloglucanase which is (a) a polypeptide produced by *Bacillus licheniformis*, ATCC 14580, or (b) a polypeptide comprising an amino acid sequence as shown in  
25 positions 30-261 of SEQ ID NO:2, or (c) an analogue of the polypeptide defined in (a) or (b) which is at least 70% homologous with said polypeptide, or is derived from said polypeptide by substitution, deletion or addition of one or several amino acids, or is immunologically reactive with a  
30 polyclonal antibody raised against said polypeptide in purified form; and to an isolated polynucleotide molecule encoding a polypeptide having xyloglucanase activity selected from (a) polynucleotide molecules comprising a nucleotide sequence as shown in SEQ ID NO:1 from nucleotide 88 to nucleotide 783; (b)  
35 polynucleotide molecules that encode a polypeptide that is at least 70% identical to the amino acid sequence of SEQ ID NO:2 from amino acid residue 30 to amino acid residue 261; and (c) degenerate nucleotide sequences of (a) or (b); and to an isolated polynucleotide molecule encoding a polypeptide having

xyloglucanase activity selected from the group consisting of:

- (a) polynucleotide molecules comprising a nucleotide sequence as shown in SEQ ID NO:3 from nucleotide 1 to nucleotide 1611;
- (b) polynucleotide molecules that encode a polypeptide that is at least 70% identical to the amino acid sequence of SEQ ID NO:4 from amino acid residue 1 to amino acid residue 537; and
- (c) degenerate nucleotide sequences of (a) or (b).

In further aspects, the invention provides an expression vector comprising a DNA segment which is eg a polynucleotide molecule of the invention; a cell comprising the DNA segment or the expression vector; and a method of producing an enzyme exhibiting xyloglucanase activity, which method comprises culturing the cell under conditions permitting the production of the enzyme, and recovering the enzyme from the culture.

In yet another aspect the invention provides an isolated enzyme exhibiting xyloglucanase activity, characterized in (i) being free from homologous impurities and (ii) the enzyme is produced by the method described above.

The novel enzyme of the present invention is useful for the treatment of cellulosic material, especially cellulose-containing fiber, yarn, woven or non-woven fabric. The treatment can be carried out during the processing of cellulosic material into a material ready for garment manufacture or fabric manufacture, e.g. in the desizing or scouring step; or during industrial or household laundering of such fabric or garment.

Accordingly, in further aspects the present invention relates to a detergent composition comprising an enzyme having substantial xyloglucanase activity in the alkaline range; and to use of the enzyme of the invention for the treatment of cellulose-containing fibers, yarn, woven or non-woven fabric.

The present invention has now made it possible to use a truly enzymatic scouring process in the preparation of cellulosic material e.g. for proper response in subsequent dyeing operations. Further, it is contemplated that detergent compositions comprising the novel enzyme are capable of removing or bleaching certain soils or stains present on laundry, especially soils and spots resulting from xyloglucan-containing food, plants, and the like. It is also contemplated that treatment with detergent compositions comprising the novel

enzyme can prevent binding of certain soils to the xyloglucan left on the cellulosic material.

#### DETAILED DESCRIPTION OF THE INVENTION

5 Cellulases are found in more than 10 different families of glycosyl hydrolases. Some of the cellulases also exhibit xyloglucanase activity. Today, such cellulases have been found among those classified in the families 5, 7 and 12. The substrate specificity is, however, not directly correlated with  
10 the family: within a family the main enzymatic activity can be cellulase or mannanase (family 5), or lichenase,  $\beta$ -1,3-glucanase or xyloglucantransferase activity (family 16). The only enzyme hitherto disclosed as having activity against xyloglucan as the major or main enzymatic activity is the *Aspergillus aculeatus* EG  
15 II disclosed in WO 94/14953. As mentioned above, this activity has not yet an entry in the official Enzyme Nomenclature.

In the present context, the term "enzyme preparation" is intended to mean either be a conventional enzymatic fermentation product, possibly isolated and purified, from a single species  
20 of a microorganism, such preparation usually comprising a number of different enzymatic activities; or a mixture of monocomponent enzymes, preferably enzymes derived from bacterial or fungal species by using conventional recombinant techniques, which enzymes have been fermented and possibly isolated and purified  
25 separately and which may originate from different species, preferably fungal or bacterial species; or the fermentation product of a microorganism which acts as a host cell for expression of a recombinant xyloglucanase, but which microorganism simultaneously produces other enzymes, e.g.  
30 xyloglucanases, proteases, or cellulases, being naturally occurring fermentation products of the microorganism, i.e. the enzyme complex conventionally produced by the corresponding naturally occurring microorganism.

In a preferred embodiment, the xyloglucanase has a relative  
35 activity at pH 7 of at least 50%, preferably at least 75%, more preferably at least 80%, especially at least 90%, compared to the activity at the optimal pH.

In another preferred embodiment, the xyloglucanase has a relative activity at pH 8 of at least 50%, preferably at least



60%, more preferably at least 75%, especially at least 90%, compared to the activity at the optimal pH.

In yet another preferred embodiment, the xyloglucanase has a relative activity at pH 9 of at least 10%, preferably at least 20%, more preferably at least 25%, compared to the activity at the optimal pH.

In yet another preferred embodiment, the xyloglucanase has a relative activity at pH 9.5 of at least 5%, preferably at least 10%, more preferably at least 15%, compared to the activity at the optimal pH.

In yet another preferred embodiment, the xyloglucanase has a relative activity at pH 10 of at least 5%, compared to the activity at the optimal pH.

In another preferred embodiment the xyloglucanase has a relative activity at a temperature of 50°C, preferably of at least 60%, preferably at least 70%, compared to the activity at the optimal temperature.

In yet another preferred embodiment, at a temperature of 60°C, the relative xyloglucanase activity is at least 40%, preferably at least 50%; at a temperature of 70°C, the relative xyloglucanase activity is at least 40%, preferably at least 45%, especially at least 50%.

In a preferred embodiment, the xyloglucanase has a minor or no activity on cellulose or cellulose derivative substrates. A conventional substrate for determining cellulase activity (endo- $\beta$ -1,4-glucanase activity, jf. EC 3.2.1.4) is carboxymethylcellulose (CMC). Another conventional substrate for determining cellulase or cellobiohydrolase activity (EC 3.2.1.91) is Avicel which is a micro-crystalline cellulose well known by the person skilled in the art. Preferably, the ratio of maximum xyloglucanase activity to maximum activity on either CMC or Avicel is at least 2:1, more preferably at least 3:1, more preferably at least 4:1, even more preferably at least 5:1, even more preferably at least 8:1, especially at least 10:1.

The xyloglucanase preparation of the invention may further comprise one or more enzymes selected from the group consisting of proteases, cellulases (endo- $\beta$ -1,4-glucanases),  $\beta$ -glucanases (endo- $\beta$ -1,3(4)-glucanases), lipases, cutinases, peroxidases,

laccases, amylases, glucoamylases, pectinases, reductases, oxidases, phenoloxidases, ligninases, pullulanases, arabinanases, hemicellulases, mannanases, galactanases, xylanases, pectin acetyl esterases, rhamnogalacturonan acetyl  
5 esterases, polygalacturonases, rhamnogalacturonases, pectin lyases, pectate lyases, pectin methylesterases, cellobiohydrolases, transglutaminases; or mixtures thereof. In a preferred embodiment, one or more or all enzymes in the preparation is produced by using recombinant techniques, i.e.  
10 the enzyme(s) is/are mono-component enzyme(s) which is/are mixed with the other enzyme(s) to form an enzyme preparation with the desired enzyme blend.

In another aspect, the present invention also relates to a method of producing the enzyme preparation of the invention, the  
15 method comprising culturing a microorganism capable of producing the xyloglucanase under conditions permitting the production of the enzyme, and recovering the enzyme from the culture.

Culturing may be carried out using conventional fermentation techniques, e.g. culturing in shake flasks or fermentors with  
20 agitation to ensure sufficient aeration on a growth medium inducing production of the xyloglucanase enzyme. The growth medium may contain a conventional N-source such as peptone, yeast extract or casamino acids, a reduced amount of a conventional C-source such as dextrose or sucrose, and an  
25 inducer such as xyloglucan or composit plant substrates such as cereal brans (e.g. wheat bran or rice husk). The recovery may be carried out using conventional techniques, e.g. separation of bio-mass and supernatant by centrifugation or filtration, recovery of the supernatant or disruption of cells if the enzyme  
30 of interest is intracellular, perhaps followed by further purification as described in EP 0 406 314 or by crystallization as described in WO 97/15660.

In the present context the term "expression vector" denotes a DNA molecule, linear or circular, that comprises a segment  
35 encoding a polypeptide of interest operably linked to additional segments that provide for its transcription. Such additional segments may include promoter and terminator sequences, and may optionally include one or more origins of replication, one or more selectable markers, an enhancer, a polyadenylation signal,

and the like. Expression vectors are generally derived from plasmid or viral DNA, or may contain elements of both. The expression vector of the invention may be any expression vector that is conveniently subjected to recombinant DNA procedures, 5 and the choice of vector will often depend on the host cell into which the vector is to be introduced. Thus, the vector may be an autonomously replicating vector, i.e. a vector which exists as an extrachromosomal entity, the replication of which is independent of chromosomal replication, e.g. a plasmid. Alternatively, 10 the vector may be one which, when introduced into a host cell, is integrated into the host cell genome and replicated together with the chromosome(s) into which it has been integrated.

The term "recombinant expressed" or "recombinantly expressed" used herein in connection with expression of a 15 polypeptide or protein is defined according to the standard definition in the art. Recombinantly expression of a protein is generally performed by using an expression vector as described immediately above.

The term "isolated", when applied to a polynucleotide 20 molecule, denotes that the polynucleotide has been removed from its natural genetic milieu and is thus free of other extraneous or unwanted coding sequences, and is in a form suitable for use within genetically engineered protein production systems. Such isolated molecules are those that are separated from their 25 natural environment and include cDNA and genomic clones. Isolated DNA molecules of the present invention are free of other genes with which they are ordinarily associated, but may include naturally occurring 5' and 3' untranslated regions such as promoters and terminators. The identification of associated 30 regions will be evident to one of ordinary skill in the art (see for example, Dynan and Tijan, Nature 316:774-78, 1985). The term "an isolated polynucleotide" may alternatively be termed "a cloned polynucleotide".

When applied to a protein/polypeptide, the term "isolated" 35 indicates that the protein is found in a condition other than its native environment. In a preferred form, the isolated protein is substantially free of other proteins, particularly other homologous proteins (i.e. "homologous impurities" (see

below)). It is preferred to provide the protein in a greater than 40% pure form, more preferably greater than 60% pure form.

Even more preferably it is preferred to provide the protein in a highly purified form, i.e., greater than 80% pure, more  
5 preferably greater than 95% pure, and even more preferably greater than 99% pure, as determined by SDS-PAGE.

The term "isolated protein/polypeptide" may alternatively be termed "purified protein/polypeptide".

The term "homologous impurities" means any impurity (e.g.  
10 another polypeptide than the polypeptide of the invention) which originate from the homologous cell where the polypeptide of the invention is originally obtained from.

The term "obtained from" as used herein in connection with a specific microbial source, means that the polynucleotide  
15 and/or polypeptide produced by the specific source, or by a cell in which a gene from the source have been inserted.

The term "operably linked", when referring to DNA segments, denotes that the segments are arranged so that they function in concert for their intended purposes, e.g. transcription  
20 initiates in the promoter and proceeds through the coding segment to the terminator

The term "polynucleotide" denotes a single- or double-stranded polymer of deoxyribonucleotide or ribonucleotide bases read from the 5' to the 3' end. Polynucleotides include RNA and  
25 DNA, and may be isolated from natural sources, synthesized in vitro, or prepared from a combination of natural and synthetic molecules.

The term "complements of polynucleotide molecules" denotes polynucleotide molecules having a complementary base sequence  
30 and reverse orientation as compared to a reference sequence. For example, the sequence 5' ATGCACGGG 3' is complementary to 5' CCCGTGCAT 3'.

The term "degenerate nucleotide sequence" denotes a sequence of nucleotides that includes one or more degenerate codons (as  
35 compared to a reference polynucleotide molecule that encodes a polypeptide). Degenerate codons contain different triplets of nucleotides, but encode the same amino acid residue (i.e., GAU and GAC triplets each encode Asp).

The term "promoter" denotes a portion of a gene containing DNA sequences that provide for the binding of RNA polymerase and initiation of transcription. Promoter sequences are commonly, but not always, found in the 5' non-coding regions of genes.

5 The term "secretory signal sequence" denotes a DNA sequence that encodes a polypeptide (a "secretory peptide") that, as a component of a larger polypeptide, directs the larger polypeptide through a secretory pathway of a cell in which it is synthesized. The larger peptide is commonly cleaved to remove  
10 the secretory peptide during transit through the secretory pathway.

#### POLYNUCLEOTIDES:

Within preferred embodiments of the invention an isolated  
15 polynucleotide of the invention will hybridize to similar sized regions of SEQ ID NO:1 or SEQ ID NO:3, or a sequence complementary thereto, under at least medium stringency conditions.

In particular polynucleotides of the invention will  
20 hybridize to a denatured double-stranded DNA probe comprising either the full sequence shown in SEQ ID NO:3 or the full sequence shown in positions 88-783 of SEQ ID NO:1 or any probe comprising a subsequence of SEQ ID NO:3 or SEQ ID NO:1 having a length of at least about 100 base pairs under at least medium  
25 stringency conditions, but preferably at high stringency conditions as described in detail below. Suitable experimental conditions for determining hybridization at medium, or high stringency between a nucleotide probe and a homologous DNA or RNA sequence involves presoaking of the filter containing the  
30 DNA fragments or RNA to hybridize in 5 x SSC (Sodium chloride/Sodium citrate, Sambrook et al. 1989) for 10 min, and prehybridization of the filter in a solution of 5 x SSC, 5 x Denhardt's solution (Sambrook et al. 1989), 0.5 % SDS and 100 µg/ml of denatured sonicated salmon sperm DNA (Sambrook et al.  
35 1989), followed by hybridization in the same solution containing a concentration of 10ng/ml of a random-primed (Feinberg, A. P. and Vogelstein, B. (1983) Anal. Biochem. 132:6-13), 32P-dCTP-labeled (specific activity higher than 1 x 10<sup>9</sup> cpm/µg ) probe for 12 hours at ca. 45°C. The filter is then washed twice for 30

minutes in 2 x SSC, 0.5 % SDS at least 60°C (medium stringency), still more preferably at least 65°C (medium/high stringency), even more preferably at least 70°C (high stringency), and even more preferably at least 75°C (very high stringency).

5 Molecules to which the oligonucleotide probe hybridizes under these conditions are detected using a x-ray film.

As previously noted, the isolated polynucleotides of the present invention include DNA and RNA. Methods for isolating DNA and RNA are well known in the art. DNA and RNA encoding  
10 genes of interest can be cloned in Gene Banks or DNA libraries by means of methods known in the art.

Polynucleotides encoding polypeptides having endogucanase activity of the invention are then identified and isolated by, for example, hybridization or PCR.

15 The present invention further provides counterpart polypeptides and polynucleotides from different bacterial strains (orthologs or paralogs). Of particular interest are xyloglucanase polypeptides from gram-positive alkalophilic strains, including species of *Bacillus*.

20 Species homologues of a polypeptide with xyloglucanase activity of the invention can be cloned using information and compositions provided by the present invention in combination with conventional cloning techniques. For example, a DNA sequence of the present invention can be cloned using  
25 chromosomal DNA obtained from a cell type that expresses the protein. Suitable sources of DNA can be identified by probing Northern blots with probes designed from the sequences disclosed herein. A library is then prepared from chromosomal DNA of a positive cell line. A DNA sequence of the invention encoding an  
30 polypeptide having xyloglucanase activity can then be isolated by a variety of methods, such as by probing with probes designed from the sequences disclosed in the present specification and claims or with one or more sets of degenerate probes based on the disclosed sequences. A DNA sequence of the invention can  
35 also be cloned using the polymerase chain reaction, or PCR (Mullis, U.S. Patent 4,683,202), using primers designed from the sequences disclosed herein. Within an additional method, the DNA library can be used to transform or transfect host cells, and expression of the DNA of interest can be detected with an

antibody (monoclonal or polyclonal) raised against the xyloglucanase cloned from *B. licheniformis*, ATCC 14580, or from *B. agaradhaerens*, NCIMB 40482, expressed and purified as described in Materials and Methods and Examples 5, 6 and 7, or  
5 by an activity test relating to a polypeptide having xyloglucanase activity.

#### POLYPEPTIDES:

The sequence of SEQ ID NO:4 and of amino acids nos. 30-261  
10 of SEQ ID NO: 2, respectively, is a mature xyloglucanase sequence of the catalytic active domain. The present invention also provides xyloglucanase polypeptides that are substantially homologous to the polypeptide of SEQ ID NO:2 or SEQ ID NO:4 and species homologs (paralogs or orthologs) thereof. The term  
15 "substantially homologous" is used herein to denote polypeptides having 75%, preferably at least 80%, more preferably at least 85%, and even more preferably at least 90%, sequence identity to the sequence shown in SEQ ID NO:4 or in amino acids nos. 30-261 of SEQ ID NO:2 or their orthologs or paralogs. Such polypeptides  
20 will more preferably be at least 95% identical, and most preferably 98% or more identical to the sequence shown in SEQ ID NO:4 or in amino acids nos. 226-490 of SEQ ID NO:2 or its orthologs or paralogs. Percent sequence identity is determined by conventional methods, by means of computer programs known in  
25 the art such as GAP provided in the GCG program package (Program Manual for the Wisconsin Package, Version 8, August 1994, Genetics Computer Group, 575 Science Drive, Madison, Wisconsin, USA 53711) as disclosed in Needleman, S.B. and Wunsch, C.D., (1970), Journal of Molecular Biology, 48, 443-453, which is  
30 hereby incorporated by reference in its entirety. GAP is used with the following settings for polypeptide sequence comparison: GAP creation penalty of 3.0 and GAP extension penalty of 0.1.

Sequence identity of polynucleotide molecules is determined by similar methods using GAP with the following settings for DNA  
35 sequence comparison: GAP creation penalty of 5.0 and GAP extension penalty of 0.3.

Substantially homologous proteins and polypeptides are characterized as having one or more amino acid substitutions, deletions or additions. These changes are preferably of a minor

nature, that is conservative amino acid substitutions (see Table 2) and other substitutions that do not significantly affect the folding or activity of the protein or polypeptide; small deletions, typically of one to about 30 amino acids; and small amino- or carboxyl-terminal extensions, such as an amino-terminal methionine residue, a small linker peptide of up to about 20-25 residues, or a small extension that facilitates purification (an affinity tag), such as a poly-histidine tract, protein A (Nilsson et al., *EMBO J.* 4:1075, 1985; Nilsson et al., *Methods Enzymol.* 198:3, 1991. See, in general Ford et al., *Protein Expression and Purification* 2: 95-107, 1991, which is incorporated herein by reference. DNAs encoding affinity tags are available from commercial suppliers (e.g., Pharmacia Biotech, Piscataway, NJ; New England Biolabs, Beverly, MA).

However, even though the changes described above preferably are of a minor nature, such changes may also be of a larger nature such as fusion of larger polypeptides of up to 300 amino acids or more both as amino- or carboxyl-terminal extensions to a polypeptide of the invention having xyloglucanase activity.

Table 1

Conservative amino acid substitutions

25	Basic:	arginine
		lysine
		histidine
	Acidic:	glutamic acid
		aspartic acid
	Polar:	glutamine
		asparagine
30	Hydrophobic:	leucine
		isoleucine
		valine
35	Aromatic:	phenylalanine
		tryptophan
		tyrosine
	Small:	glycine
		alanine
		serine
		threonine



## methionine

In addition to the 20 standard amino acids, non-standard amino acids (such as 4-hydroxyproline, 6-N-methyl lysine, 2-aminoisobutyric acid, isovaline and α-methyl serine) may be substituted for amino acid residues of a polypeptide according to the invention. A limited number of non-conservative amino acids, amino acids that are not encoded by the genetic code, and unnatural amino acids may be substituted for amino acid residues. "Unnatural amino acids" have been modified after protein synthesis, and/or have a chemical structure in their side chain(s) different from that of the standard amino acids. Unnatural amino acids can be chemically synthesized, or preferably, are commercially available, and include pipecolic acid, thiazolidine carboxylic acid, dehydroproline, 3- and 4-methylproline, and 3,3-dimethylproline.

Essential amino acids in the xyloglucanase polypeptides of the present invention can be identified according to procedures known in the art, such as site-directed mutagenesis or alanine-scanning mutagenesis (Cunningham and Wells, Science 244: 1081-1085, 1989). In the latter technique, single alanine mutations are introduced at every residue in the molecule, and the resultant mutant molecules are tested for biological activity (i.e. xyloglucanase activity) to identify amino acid residues that are critical to the activity of the molecule. See also, Hilton et al., J. Biol. Chem. 271:4699-4708, 1996. The active site of the enzyme or other biological interaction can also be determined by physical analysis of structure, as determined by such techniques as nuclear magnetic resonance, crystallography, electron diffraction or photoaffinity labeling, in conjunction with mutation of putative contact site amino acids. See, for example, de Vos et al., Science 255:306-312, 1992; Smith et al., J. Mol. Biol. 224:899-904, 1992; Wlodaver et al., FEBS Lett. 309:59-64, 1992. The identities of essential amino acids can also be inferred from analysis of homologies with polypeptides which are related to a polypeptide according to the invention.

Multiple amino acid substitutions can be made and tested using known methods of mutagenesis, recombination and/or shuffling followed by a relevant screening procedure, such as

those disclosed by Reidhaar-Olson and Sauer (Science 241:53-57, 1988), Bowie and Sauer (Proc. Natl. Acad. Sci. USA 86:2152-2156, 1989), WO95/17413, or WO 95/22625. Briefly, these authors disclose methods for simultaneously randomizing two or more  
5 positions in a polypeptide, or recombination/shuffling of different mutations (WO95/17413, WO95/22625), followed by selecting for functional a polypeptide, and then sequencing the mutagenized polypeptides to determine the spectrum of allowable substitutions at each position. Other methods that can be used  
10 include phage display (e.g., Lowman et al., Biochem. 30:10832-10837, 1991; Ladner et al., U.S. Patent No. 5,223,409; Huse, WIPO Publication WO 92/06204) and region-directed mutagenesis (Derbyshire et al., Gene 46:145, 1986; Ner et al., DNA 7:127, 1988).

15 Mutagenesis/shuffling methods as disclosed above can be combined with high-throughput, automated screening methods to detect activity of cloned, mutagenized polypeptides in host cells. Mutagenized DNA molecules that encode active polypeptides can be recovered from the host cells and rapidly sequenced using  
20 modern equipment. These methods allow the rapid determination of the importance of individual amino acid residues in a polypeptide of interest, and can be applied to polypeptides of unknown structure.

Using the methods discussed above, one of ordinary skill in  
25 the art can identify and/or prepare a variety of polypeptides that are substantially homologous to SEQ ID NO:4 or residues 30 to 261 of SEQ ID NO: 2 and retain the xyloglucanase activity of the wild-type protein.

The xyloglucanase enzyme of the invention may, in addition  
30 to the enzyme core comprising the catalytically domain, also comprise a cellulose binding domain (CBD), the cellulose binding domain and enzyme core (the catalytically active domain) of the enzyme being operably linked. The cellulose binding domain (CBD) may exist as an integral part the encoded enzyme, or a CBD from  
35 another origin may be introduced into the xyloglucanase thus creating an enzyme hybride. In this context, the term "cellulose-binding domain" is intended to be understood as defined by Peter Tomme et al. "Cellulose-Binding Domains: Classification and Properties" in "Enzymatic Degradation of

Insoluble Carbohydrates", John N. Saddler and Michael H. Penner (Eds.), ACS Symposium Series, No. 618, 1996. This definition classifies more than 120 cellulose-binding domains into 10 families (I-X), and demonstrates that CBDs are found in various enzymes such as cellulases, xylanases, mannanases, arabinofuranosidases, acetyl esterases and chitinases. CBDs have also been found in algae, e.g. the red alga *Porphyra purpurea* as a non-hydrolytic polysaccharide-binding protein, see Tomme et al., *op.cit.* However, most of the CBDs are from cellulases and xylanases, CBDs are found at the N and C termini of proteins or are internal. Enzyme hybrids are known in the art, see e.g. WO 90/00609 and WO 95/16782, and may be prepared by transforming into a host cell a DNA construct comprising at least a fragment of DNA encoding the cellulose-binding domain ligated, with or without a linker, to a DNA sequence encoding the xyloglucanase and growing the host cell to express the fused gene. Enzyme hybrids may be described by the following formula:

CBD - MR - X

wherein CBD is the N-terminal or the C-terminal region of an amino acid sequence corresponding to at least the cellulose-binding domain; MR is the middle region (the linker), and may be a bond, or a short linking group preferably of from about 2 to about 100 carbon atoms, more preferably of from 2 to 40 carbon atoms; or is preferably from about 2 to to about 100 amino acids, more preferably of from 2 to 40 amino acids; and X is an N-terminal or C-terminal region of a polypeptide encoded by the polynucleotide molecule of the invention.

#### The xyloglucan substrate

In addition to the aforesaid about xyloglucan it should be noted that xyloglucan from tamarind seeds supplied by Megazyme, Australia has a complex branched structure with glucose, xylose, galactose and arabinose in the ratio of 45:36:16:3. Accordingly, it is strongly believed that an enzyme showing catalytic activity on this xyloglucan also has catalytic activity on other xyloglucan structures from different sources (angiosperms or gymnosperms).

#### Use in the detergent industry

During washing and wearing, dyestuff from dyed fabrics or garment will conventionally bleed from the fabric which then looks faded and worn. Removal of surface fibers from the fabric will partly restore the original colours and looks of the  
5 fabric. By the term "colour clarification", as used herein, is meant the partly restoration of the initial colours of fabric or garment throughout multiple washing cycles.

The term "de-pilling" denotes removing of pills from the fabric surface.

10 The term "soaking liquor" denotes an aqueous liquor in which laundry may be immersed prior to being subjected to a conventional washing process. The soaking liquor may contain one or more ingredients conventionally used in a washing or laundering process.

15 The term "washing liquor" denotes an aqueous liquor in which laundry is subjected to a washing process, i.e. usually a combined chemical and mechanical action either manually or in a washing machine. Conventionally, the washing liquor is an aqueous solution of a powder or liquid detergent composition.

20 The term "rinsing liquor" denotes an aqueous liquor in which laundry is immersed and treated, conventionally immediately after being subjected to a washing process, in order to rinse the laundry, i.e. essentially remove the detergent solution from the laundry. The rinsing liquor may contain a  
25 fabric conditioning or softening composition.

The laundry subjected to the method of the present invention may be conventional washable laundry. Preferably, the major part of the laundry is sewn or unsewn fabrics, including knits, wovens, denims, yarns, and toweling, made from cotton, cotton  
30 blends or natural or manmade cellulosics (e.g. originating from xylan-containing cellulose fibers such as from wood pulp) or blends thereof. Examples of blends are blends of cotton or rayon/viscose with one or more companion material such as wool, synthetic fibers (e.g. polyamide fibers, acrylic fibers,  
35 polyester fibers, polyvinyl alcohol fibers, polyvinyl chloride fibers, polyvinylidene chloride fibers, polyurethane fibers, polyurea fibers, aramid fibers), and cellulose-containing fibers (e.g. rayon/viscose, ramie, flax/linen, jute, cellulose acetate fibers, lyocell).

**DETERGENT DISCLOSURE AND EXAMPLES****Surfactant system**

The detergent compositions according to the present invention comprise a surfactant system, wherein the surfactant can be selected from nonionic and/or anionic and/or cationic and/or ampholytic and/or zwitterionic and/or semi-polar surfactants.

The surfactant is typically present at a level from 0.1% to 60% by weight.

The surfactant is preferably formulated to be compatible with enzyme components present in the composition. In liquid or gel compositions the surfactant is most preferably formulated in such a way that it promotes, or at least does not degrade, the stability of any enzyme in these compositions.

Preferred systems to be used according to the present invention comprise as a surfactant one or more of the nonionic and/or anionic surfactants described herein.

Polyethylene, polypropylene, and polybutylene oxide condensates of alkyl phenols are suitable for use as the nonionic surfactant of the surfactant systems of the present invention, with the polyethylene oxide condensates being preferred. These compounds include the condensation products of alkyl phenols having an alkyl group containing from about 6 to about 14 carbon atoms, preferably from about 8 to about 14 carbon atoms, in either a straight chain or branched-chain configuration with the alkylene oxide. In a preferred embodiment, the ethylene oxide is present in an amount equal to from about 2 to about 25 moles, more preferably from about 3 to about 15 moles, of ethylene oxide per mole of alkyl phenol. Commercially available nonionic surfactants of this type include Igepal<sup>TM</sup> CO-630, marketed by the GAF Corporation; and Triton<sup>TM</sup> X-45, X-114, X-100 and X-102, all marketed by the Rohm & Haas Company. These surfactants are commonly referred to as alkylphenol alkoxylates (e.g., alkyl phenol ethoxylates).


The condensation products of primary and secondary aliphatic alcohols with about 1 to about 25 moles of ethylene oxide are suitable for use as the nonionic surfactant of the nonionic surfactant systems of the present invention. The alkyl

chain of the aliphatic alcohol can either be straight or branched, primary or secondary, and generally contains from about 8 to about 22 carbon atoms. Preferred are the condensation products of alcohols having an alkyl group containing from about 5 8 to about 20 carbon atoms, more preferably from about 10 to about 18 carbon atoms, with from about 2 to about 10 moles of ethylene oxide per mole of alcohol. About 2 to about 7 moles of ethylene oxide and most preferably from 2 to 5 moles of ethylene oxide per mole of alcohol are present in said condensation products. Examples of commercially available nonionic surfactants of this type include Tergitol<sup>TM</sup> 15-S-9 (The condensation product of C<sub>11</sub>-C<sub>15</sub> linear alcohol with 9 moles ethylene oxide), Tergitol<sup>TM</sup> 24-L-6 NMW (the condensation product of C<sub>12</sub>-C<sub>14</sub> primary alcohol with 6 moles ethylene oxide with a narrow 15 molecular weight distribution), both marketed by Union Carbide Corporation; Neodol<sup>TM</sup> 45-9 (the condensation product of C<sub>14</sub>-C<sub>15</sub> linear alcohol with 9 moles of ethylene oxide), Neodol<sup>TM</sup> 23-3 (the condensation product of C<sub>12</sub>-C<sub>13</sub> linear alcohol with 3.0 moles of ethylene oxide), Neodol<sup>TM</sup> 45-7 (the condensation 20 product of C<sub>14</sub>-C<sub>15</sub> linear alcohol with 7 moles of ethylene oxide), Neodol<sup>TM</sup> 45-5 (the condensation product of C<sub>14</sub>-C<sub>15</sub> linear alcohol with 5 moles of ethylene oxide) marketed by Shell Chemical Company, Kyro<sup>TM</sup> EOB (the condensation product of C<sub>13</sub>-C<sub>15</sub> alcohol with 9 moles ethylene oxide), marketed by The Procter & 25 Gamble Company, and Genapol LA 050 (the condensation product of C<sub>12</sub>-C<sub>14</sub> alcohol with 5 moles of ethylene oxide) marketed by Hoechst. Preferred range of HLB in these products is from 8-11 and most preferred from 8-10.

Also useful as the nonionic surfactant of the surfactant 30 systems of the present invention are alkylpolysaccharides disclosed in US 4,565,647, having a hydrophobic group containing from about 6 to about 30 carbon atoms, preferably from about 10 to about 16 carbon atoms and a polysaccharide, e.g. a polyglycoside, hydrophilic group containing from about 1.3 to 35 about 10, preferably from about 1.3 to about 3, most preferably from about 1.3 to about 2.7 saccharide units. Any reducing saccharide containing 5 or 6 carbon atoms can be used, e.g., glucose, galactose and galactosyl moieties can be substituted for the glucosyl moieties (optionally the hydrophobic group is

attached at the 2-, 3-, 4-, etc. positions thus giving a glucose or galactose as opposed to a glucoside or galactoside). The intersaccharide bonds can be, e.g., between the one position of the additional saccharide units and the 2-, 3-, 4-, and/or 6-  
5 positions on the preceding saccharide units.

The preferred alkylpolyglycosides have the formula



10 wherein  $R^2$  is selected from the group consisting of alkyl, alkylphenyl, hydroxyalkyl, hydroxyalkylphenyl, and mixtures thereof in which the alkyl groups contain from about 10 to about 18, preferably from about 12 to about 14, carbon atoms; n is 2 or 3, preferably 2; t is from 0 to about 10, preferably 0; and  
15 x is from about 1.3 to about 10, preferably from about 1.3 to about 3, most preferably from about 1.3 to about 2.7. The glycosyl is preferably derived from glucose. To prepare these compounds, the alcohol or alkylpolyethoxy alcohol is formed first and then reacted with glucose, or a source of glucose, to  
20 form the glucoside (attachment at the 1-position). The additional glycosyl units can then be attached between their 1-position and the preceding glycosyl units 2-, 3-, 4-, and/or 6-position, preferably predominantly the 2-position.

The condensation products of ethylene oxide with a  
25 hydrophobic base formed by the condensation of propylene oxide with propylene glycol are also suitable for use as the additional nonionic surfactant systems of the present invention. The hydrophobic portion of these compounds will preferably have a molecular weight from about 1500 to about 1800 and will  
30 exhibit water insolubility. The addition of polyoxyethylene moieties to this hydrophobic portion tends to increase the water solubility of the molecule as a whole, and the liquid character of the product is retained up to the point where the polyoxyethylene content is about 50% of the total weight of the  
35 condensation product, which corresponds to condensation with up to about 40 moles of ethylene oxide. Examples of compounds of this type include certain of the commercially available Pluronic<sup>TM</sup> surfactants, marketed by BASF.

Also suitable for use as the nonionic surfactant of the

nonionic surfactant system of the present invention, are the condensation products of ethylene oxide with the product resulting from the reaction of propylene oxide and ethylenediamine. The hydrophobic moiety of these products  
 5 consists of the reaction product of ethylenediamine and excess propylene oxide, and generally has a molecular weight of from about 2500 to about 3000. This hydrophobic moiety is condensed with ethylene oxide to the extent that the condensation product contains from about 40% to about 80% by weight of  
 10 polyoxyethylene and has a molecular weight of from about 5,000 to about 11,000. Examples of this type of nonionic surfactant include certain of the commercially available Tetronic<sup>TM</sup> compounds, marketed by BASF.

Preferred for use as the nonionic surfactant of the  
 15 surfactant systems of the present invention are polyethylene oxide condensates of alkyl phenols, condensation products of primary and secondary aliphatic alcohols with from about 1 to about 25 moles of ethyleneoxide, alkylpolysaccharides, and mixtures hereof. Most preferred are C<sub>8</sub>-C<sub>14</sub> alkyl phenol  
 20 ethoxylates having from 3 to 15 ethoxy groups and C<sub>8</sub>-C<sub>18</sub> alcohol ethoxylates (preferably C<sub>10</sub> avg.) having from 2 to 10 ethoxy groups, and mixtures thereof.  
 Highly preferred nonionic surfactants are polyhydroxy fatty acid amide surfactants of the formula




wherein R<sup>1</sup> is H, or R<sup>1</sup> is C<sub>1-4</sub> hydrocarbyl, 2-hydroxyethyl, 2-hydroxypropyl or a mixture thereof, R<sup>2</sup> is C<sub>5-31</sub> hydrocarbyl, and  
 30 Z is a polyhydroxyhydrocarbyl having a linear hydrocarbyl chain with at least 3 hydroxyls directly connected to the chain, or an alkoxylated derivative thereof. Preferably, R<sup>1</sup> is methyl, R<sup>2</sup> is straight C<sub>11-15</sub> alkyl or C<sub>16-18</sub> alkyl or alkenyl chain such as coconut alkyl or mixtures thereof, and Z is derived from a  
 35 reducing sugar such as glucose, fructose, maltose or lactose, in a reductive amination reaction.

Highly preferred anionic surfactants include alkyl alkoxylated sulfate surfactants. Examples hereof are water soluble salts or acids of the formula RO(A)<sub>m</sub>SO<sub>3</sub>M wherein R is an



unsubstituted C<sub>10</sub>-C<sub>24</sub> alkyl or hydroxyalkyl group having a C<sub>10</sub>-C<sub>24</sub> alkyl component, preferably a C<sub>12</sub>-C<sub>20</sub> alkyl or hydro-xyalkyl, more preferably C<sub>12</sub>-C<sub>18</sub> alkyl or hydroxyalkyl, A is an ethoxy or propoxy unit, m is greater than zero, typically between about 0.5 and about 6, more preferably between about 0.5 and about 3, and M is H or a cation which can be, for example, a metal cation (e.g., sodium, potassium, lithium, calcium, magnesium, etc.), ammonium or substituted-ammonium cation. Alkyl ethoxylated sulfates as well as alkyl propoxylated sulfates are contemplated herein. Specific examples of substituted ammonium cations include methyl-, dimethyl, trimethyl-ammonium cations and quaternary ammonium cations such as tetramethyl-ammonium and dimethyl piperdinium cations and those derived from alkylamines such as ethylamine, diethylamine, triethylamine, mixtures thereof, and the like. Exemplary surfactants are C<sub>12</sub>-C<sub>18</sub> alkyl polyethoxylate (1.0) sulfate (C<sub>12</sub>-C<sub>18</sub>E(1.0)M), C<sub>12</sub>-C<sub>18</sub> alkyl polyethoxylate (2.25) sulfate (C<sub>12</sub>-C<sub>18</sub>(2.25)M, and C<sub>12</sub>-C<sub>18</sub> alkyl polyethoxylate (3.0) sulfate (C<sub>12</sub>-C<sub>18</sub>E(3.0)M), and C<sub>12</sub>-C<sub>18</sub> alkyl polyethoxylate (4.0) sulfate (C<sub>12</sub>-C<sub>18</sub>E(4.0)M), wherein M is conveniently selected from sodium and potassium. Suitable anionic surfactants to be used are alkyl ester sulfonate surfactants including linear esters of C<sub>8</sub>-C<sub>20</sub> carboxylic acids (i.e., fatty acids) which are sulfonated with gaseous SO<sub>3</sub> according to "The Journal of the American Oil Chemists Society", 52 (1975), pp. 323-329. Suitable starting materials would include natural fatty substances as derived from tallow, palm oil, etc.

The preferred alkyl ester sulfonate surfactant, especially for laundry applications, comprise alkyl ester sulfonate surfactants of the structural formula:



wherein R<sup>3</sup> is a C<sub>8</sub>-C<sub>20</sub> hydrocarbyl, preferably an alkyl, or combination thereof, R<sup>4</sup> is a C<sub>1</sub>-C<sub>6</sub> hydrocarbyl, preferably an alkyl, or combination thereof, and M is a cation which forms a

water soluble salt with the alkyl ester sulfonate. Suitable salt-forming cations include metals such as sodium, potassium, and lithium, and substituted or unsubstituted ammonium cations, such as monoethanolamine, diethanolamine, and triethanolamine. Preferably,  $R^3$  is  $C_{10}$ - $C_{16}$  alkyl, and  $R^4$  is methyl, ethyl or isopropyl. Especially preferred are the methyl ester sulfonates wherein  $R^3$  is  $C_{10}$ - $C_{16}$  alkyl.

Other suitable anionic surfactants include the alkyl sulfate surfactants which are water soluble salts or acids of the formula  $ROSO_3M$  wherein R preferably is a  $C_{10}$ - $C_{24}$  hydrocarbyl, preferably an alkyl or hydroxyalkyl having a  $C_{10}$ - $C_{20}$  alkyl component, more preferably a  $C_{12}$ - $C_{18}$  alkyl or hydroxyalkyl, and M is H or a cation, e.g., an alkali metal cation (e.g. sodium, potassium, lithium), or ammonium or substituted ammonium (e.g. methyl-, dimethyl-, and trimethyl ammonium cations and quaternary ammonium cations such as tetramethyl-ammonium and dimethyl piperdinium cations and quaternary ammonium cations derived from alkylamines such as ethylamine, diethylamine, triethylamine, and mixtures thereof, and the like). Typically, alkyl chains of  $C_{12}$ - $C_{16}$  are preferred for lower wash temperatures (e.g. below about  $50^\circ\text{C}$ ) and  $C_{16}$ - $C_{18}$  alkyl chains are preferred for higher wash temperatures (e.g. above about  $50^\circ\text{C}$ ).

Other anionic surfactants useful for deterative purposes can also be included in the laundry detergent compositions of the present invention. These can include salts (including, for example, sodium, potassium, ammonium, and substituted ammonium salts such as mono- di- and triethanolamine salts) of soap,  $C_8$ - $C_{22}$  primary or secondary alkanesulfonates,  $C_8$ - $C_{24}$  olefinsulfonates, sulfonated polycarboxylic acids prepared by sulfonation of the pyrolyzed product of alkaline earth metal citrates, e.g., as described in British patent specification No. 1,082,179,  $C_8$ - $C_{24}$  alkylpolyglycoethersulfates (containing up to 10 moles of ethylene oxide); alkyl glycerol sulfonates, fatty acyl glycerol sulfonates, fatty oleyl glycerol sulfates, alkyl phenol ethylene oxide ether sulfates, paraffin sulfonates, alkyl phosphates, isethionates such as the acyl isethionates, N-acyl taurates, alkyl succinamates and sulfosuccinates, monoesters of sulfosuccinates (especially saturated and unsaturated  $C_{12}$ - $C_{18}$

monoesters) and diesters of sulfosuccinates (especially saturated and unsaturated C<sub>6</sub>-C<sub>12</sub> diesters), acyl sarcosinates, sulfates of alkylpolysaccharides such as the sulfates of alkylpolyglucoside (the nonionic nonsulfated compounds being described below), branched primary alkyl sulfates, and alkyl polyethoxy carboxylates such as those of the formula  $RO(CH_2CH_2O)_k-CH_2COO-M^+$  wherein R is a C<sub>8</sub>-C<sub>22</sub> alkyl, k is an integer from 1 to 10, and M is a soluble salt forming cation. Resin acids and hydrogenated resin acids are also suitable, such as rosin, hydrogenated rosin, and resin acids and hydrogenated resin acids present in or derived from tall oil.


Alkylbenzene sulfonates are highly preferred. Especially preferred are linear (straight-chain) alkyl benzene sulfonates (LAS) wherein the alkyl group preferably contains from 10 to 18 carbon atoms.

Further examples are described in "Surface Active Agents and Detergents" (Vol. I and II by Schwartz, Perry and Berch). A variety of such surfactants are also generally disclosed in US 3,929,678, (Column 23, line 58 through Column 29, line 23, herein incorporated by reference).

When included therein, the laundry detergent compositions of the present invention typically comprise from about 1% to about 40%, preferably from about 3% to about 20% by weight of such anionic surfactants.


The laundry detergent compositions of the present invention may also contain cationic, ampholytic, zwitterionic, and semi-polar surfactants, as well as the nonionic and/or anionic surfactants other than those already described herein.

Cationic deterative surfactants suitable for use in the laundry detergent compositions of the present invention are those having one long-chain hydrocarbyl group. Examples of such cationic surfactants include the ammonium surfactants such as alkyltrimethylammonium halogenides, and those surfactants having the formula:



wherein  $R^2$  is an alkyl or alkyl benzyl group having from about 8 to about 18 carbon atoms in the alkyl chain, each  $R^3$  is selected from the group consisting of  $-\text{CH}_2\text{CH}_2-$ ,  $-\text{CH}_2\text{CH}(\text{CH}_3)-$ ,  $-\text{CH}_2\text{CH}(\text{CH}_2\text{OH})-$ ,  $-\text{CH}_2\text{CH}_2\text{CH}_2-$ , and mixtures thereof; each  $R^4$  is  
 5 selected from the group consisting of  $\text{C}_1$ - $\text{C}_4$  alkyl,  $\text{C}_1$ - $\text{C}_4$  hydroxyalkyl, benzyl ring structures formed by joining the two  $R^4$  groups,  $-\text{CH}_2\text{CHOHCHOHCOR}^6\text{CHOHCH}_2\text{OH}$ , wherein  $R^6$  is any hexose or hexose polymer having a molecular weight less than about 1000, and hydrogen when  $y$  is not 0;  $R^5$  is the same as  $R^4$  or is an  
 10 alkyl chain, wherein the total number of carbon atoms or  $R^2$  plus  $R^5$  is not more than about 18; each  $y$  is from 0 to about 10, and the sum of the  $y$  values is from 0 to about 15; and  $X$  is any compatible anion.

Highly preferred cationic surfactants are the water  
 15 soluble quaternary ammonium compounds useful in the present composition having the formula:



20 wherein  $R_1$  is  $\text{C}_8$ - $\text{C}_{16}$  alkyl, each of  $R_2$ ,  $R_3$  and  $R_4$  is independently  $\text{C}_1$ - $\text{C}_4$  alkyl,  $\text{C}_1$ - $\text{C}_4$  hydroxy alkyl, benzyl, and  $-(\text{C}_2\text{H}_4\text{O})_x\text{H}$  where  $x$  has a value from 2 to 5, and  $X$  is an anion. Not more than one of  $R_2$ ,  $R_3$  or  $R_4$  should be benzyl.


The preferred alkyl chain length for  $R_1$  is  $\text{C}_{12}$ - $\text{C}_{15}$ ,  
 25 particularly where the alkyl group is a mixture of chain lengths derived from coconut or palm kernel fat or is derived synthetically by olefin build up or OXO alcohols synthesis.

Preferred groups for  $R_2R_3$  and  $R_4$  are methyl and hydroxyethyl groups and the anion  $X$  may be selected from halide,  
 30 methosulphate, acetate and phosphate ions.

Examples of suitable quaternary ammonium compounds of formulae (i) for use herein are:

coconut trimethyl ammonium chloride or bromide;  
 coconut methyl dihydroxyethyl ammonium chloride or bromide;  
 35 decyl triethyl ammonium chloride;  
 decyl dimethyl hydroxyethyl ammonium chloride or bromide;  
 $\text{C}_{12-15}$  dimethyl hydroxyethyl ammonium chloride or bromide;  
 coconut dimethyl hydroxyethyl ammonium chloride or bromide;  
 myristyl trimethyl ammonium methyl sulphate;

lauryl dimethyl benzyl ammonium chloride or bromide;  
 lauryl dimethyl (ethenoxy)<sub>4</sub> ammonium chloride or bromide;  
 choline esters (compounds of formula (i) wherein R<sub>1</sub> is



di-alkyl imidazolines [compounds of formula (i)].

10           Other cationic surfactants useful herein are also  
 described in US 4,228,044 and in EP 000 224.

When included therein, the laundry detergent compositions  
 of the present invention typically comprise from 0.2% to about  
 25%, preferably from about 1% to about 8% by weight of such  
 15 cationic surfactants.

Ampholytic surfactants are also suitable for use in the  
 laundry detergent compositions of the present invention. These  
 surfactants can be broadly described as aliphatic derivatives of  
 secondary or tertiary amines, or aliphatic derivatives of  
 20 heterocyclic secondary and tertiary amines in which the  
 aliphatic radical can be straight- or branched-chain. One of the  
 aliphatic substituents contains at least about 8 carbon atoms,  
 typically from about 8 to about 18 carbon atoms, and at least  
 one contains an anionic water-solubilizing group, e.g. carboxy,  
 25 sulfonate, sulfate. See US 3,929,678 (column 19, lines 18-35)  
 for examples of ampholytic surfactants.

When included therein, the laundry detergent compositions  
 of the present invention typically comprise from 0.2% to about  
 15%, preferably from about 1% to about 10% by weight of such  
 30 ampholytic surfactants.

Zwitterionic surfactants are also suitable for use in  
 laundry detergent compositions. These surfactants can be broadly  
 described as derivatives of secondary and tertiary amines,  
 derivatives of heterocyclic secondary and tertiary amines, or  
 35 derivatives of quaternary ammonium, quaternary phosphonium or  
 tertiary sulfonium compounds. See US 3,929,678 (column 19, line  
 38 through column 22, line 48) for examples of zwitterionic  
 surfactants.


When included therein, the laundry detergent compositions

of the present invention typically comprise from 0.2% to about 15%, preferably from about 1% to about 10% by weight of such zwitterionic surfactants.

Semi-polar nonionic surfactants are a special category of nonionic surfactants which include water-soluble amine oxides containing one alkyl moiety of from about 10 to about 18 carbon atoms and 2 moieties selected from the group consisting of alkyl groups and hydroxyalkyl groups containing from about 1 to about 3 carbon atoms; watersoluble phosphine oxides containing one alkyl moiety of from about 10 to about 18 carbon atoms and 2 moieties selected from the group consisting of alkyl groups and hydroxyalkyl groups containing from about 1 to about 3 carbon atoms; and water-soluble sulfoxides containing one alkyl moiety from about 10 to about 18 carbon atoms and a moiety selected from the group consisting of alkyl and hydroxyalkyl moieties of from about 1 to about 3 carbon atoms.

Semi-polar nonionic detergent surfactants include the amine oxide surfactants having the formula:

20



wherein  $\text{R}^3$  is an alkyl, hydroxyalkyl, or alkyl phenyl group or mixtures thereof containing from about 8 to about 22 carbon atoms;  $\text{R}^4$  is an alkylene or hydroxyalkylene group containing from about 2 to about 3 carbon atoms or mixtures thereof;  $x$  is from 0 to about 3; and each  $\text{R}^5$  is an alkyl or hydroxyalkyl group containing from about 1 to about 3 carbon atoms or a polyethylene oxide group containing from about 1 to about 3 ethylene oxide groups. The  $\text{R}^5$  groups can be attached to each other, e.g., through an oxygen or nitrogen atom, to form a ring structure.

These amine oxide surfactants in particular include  $\text{C}_{10}$ - $\text{C}_{18}$  alkyl dimethyl amine oxides and  $\text{C}_8$ - $\text{C}_{12}$  alkoxy ethyl dihydroxy ethyl amine oxides.

When included therein, the laundry detergent compositions of the present invention typically comprise from 0.2% to about 15%, preferably from about 1% to about 10% by weight of such

semi-polar nonionic surfactants.

#### Builder system

The compositions according to the present invention may  
5 further comprise a builder system. Any conventional builder  
system is suitable for use herein including aluminosilicate  
materials, silicates, polycarboxylates and fatty acids,  
materials such as ethylenediamine tetraacetate, metal ion  
sequestrants such as aminopolyphosphonates, particularly  
10 ethylenediamine tetramethylene phosphonic acid and diethylene  
triamine pentamethylenephosphonic acid. Though less preferred  
for obvious environmental reasons, phosphate builders can also  
be used herein.

Suitable builders can be an inorganic ion exchange  
15 material, commonly an inorganic hydrated aluminosilicate  
material, more particularly a hydrated synthetic zeolite such as  
hydrated zeolite A, X, B, HS or MAP.

Another suitable inorganic builder material is layered  
silicate, e.g. SKS-6 (Hoechst). SKS-6 is a crystalline layered  
20 silicate consisting of sodium silicate ( $\text{Na}_2\text{Si}_2\text{O}_5$ ).

Suitable polycarboxylates containing one carboxy group  
include lactic acid, glycolic acid and ether derivatives thereof  
as disclosed in Belgian Patent Nos. 831,368, 821,369 and  
821,370. Polycarboxylates containing two carboxy groups include  
25 the water-soluble salts of succinic acid, malonic acid,  
(ethylenedioxy) diacetic acid, maleic acid, diglycollic acid,  
tartaric acid, tartronic acid and fumaric acid, as well as the  
ether carboxylates described in German Offenle-enschrift  
2,446,686, and 2,446,487, US 3,935,257 and the sulfinyl  
30 carboxylates described in Belgian Patent No. 840,623.  
Polycarboxylates containing three carboxy groups include, in  
particular, water-soluble citrates, aconitrates and citraconates  
as well as succinate derivatives such as the  
carboxymethyloxysuccinates described in British Patent No.  
35 1,379,241, lactoxysuccinates described in Netherlands  
Application 7205873, and the oxypolycarboxylate materials such  
as 2-oxa-1,1,3-propane tricarboxylates described in British  
Patent No. 1,387,447.

Polycarboxylates containing four carboxy groups include

oxydisuccinates disclosed in British Patent No. 1,261,829, 1,1,2,2,-ethane tetracarboxylates, 1,1,3,3-propane tetracarboxylates containing sulfo substituents include the sulfosuccinate derivatives disclosed in British Patent Nos. 5 1,398,421 and 1,398,422 and in US 3,936,448, and the sulfonated pyrolysed citrates described in British Patent No. 1,082,179, while polycarboxylates containing phosphone substituents are disclosed in British Patent No. 1,439,000.

Alicyclic and heterocyclic polycarboxylates include 10 cyclopentane-cis,cis-cis-tetracarboxylates, cyclopentadienide pentacarboxylates, 2,3,4,5-tetrahydro-furan - cis, cis, cis-tetracarboxylates, 2,5-tetrahydro-furan-cis, dicarboxylates, 2,2,5,5,-tetrahydrofuran - tetracarboxylates, 1,2,3,4,5,6-hexane - hexacarboxylates and carboxymethyl derivatives of polyhydric 15 alcohols such as sorbitol, mannitol and xylitol. Aromatic polycarboxylates include mellitic acid, pyromellitic acid and the phthalic acid derivatives disclosed in British Patent No. 1,425,343.

Of the above, the preferred polycarboxylates are hydroxy- 20 carboxylates containing up to three carboxy groups per molecule, more particularly citrates.

Preferred builder systems for use in the present compositions include a mixture of a water-insoluble aluminosilicate builder such as zeolite A or of a layered 25 silicate (SKS-6), and a water-soluble carboxylate chelating agent such as citric acid.

A suitable chelant for inclusion in the detergent compositions in accordance with the invention is ethylenediamine-N,N'-disuccinic acid (EDDS) or the alkali metal, alkaline earth 30 metal, ammonium, or substituted ammonium salts thereof, or mixtures thereof. Preferred EDDS compounds are the free acid form and the sodium or magnesium salt thereof. Examples of such preferred sodium salts of EDDS include Na<sub>2</sub>EDDS and Na<sub>4</sub>EDDS. Examples of such preferred magnesium salts of EDDS include 35 MgEDDS and Mg<sub>2</sub>EDDS. The magnesium salts are the most preferred for inclusion in compositions in accordance with the invention.

Preferred builder systems include a mixture of a water-insoluble aluminosilicate builder such as zeolite A, and a water soluble carboxylate chelating agent such as citric acid.



Other builder materials that can form part of the builder system for use in granular compositions include inorganic materials such as alkali metal carbonates, bicarbonates, silicates, and organic materials such as the organic  
5 phosphonates, amino polyalkylene phosphonates and amino polycarboxylates.

Other suitable water-soluble organic salts are the homo- or co-polymeric acids or their salts, in which the polycarboxylic acid comprises at least two carboxyl radicals  
10 separated from each other by not more than two carbon atoms.

Polymers of this type are disclosed in GB-A-1,596,756. Examples of such salts are polyacrylates of MW 2000-5000 and their copolymers with maleic anhydride, such copolymers having a molecular weight of from 20,000 to 70,000, especially about  
15 40,000.

Detergency builder salts are normally included in amounts of from 5% to 80% by weight of the composition. Preferred levels of builder for liquid detergents are from 5% to 30%.

## 20 Enzymes

Preferred detergent compositions, in addition to the enzyme preparation of the invention, comprise other enzyme(s) which provides cleaning performance and/or fabric care benefits.

Such enzymes include proteases, lipases, cutinases,  
25 amylases, cellulases, peroxidases, oxidases (e.g. laccases).

Proteases: Any protease suitable for use in alkaline solutions can be used. Suitable proteases include those of animal, vegetable or microbial origin. Microbial origin is preferred. Chemically or genetically modified mutants are included. The  
30 protease may be a serine protease, preferably an alkaline microbial protease or a trypsin-like protease. Examples of alkaline proteases are subtilisins, especially those derived from Bacillus, e.g., subtilisin Novo, subtilisin Carlsberg, subtilisin 309, subtilisin 147 and subtilisin 168 (described in  
35 WO 89/06279). Examples of trypsin-like proteases are trypsin (e.g. of porcine or bovine origin) and the Fusarium protease described in WO 89/06270.

Preferred commercially available protease enzymes include those sold under the trade names Alcalase, Savinase, Primase,

Durazym, and Esperase by Novo Nordisk A/S (Denmark), those sold under the tradename Maxatase, Maxacal, Maxapem, Properase, Purafect and Purafect OXP by Genencor International, and those sold under the tradename Opticlean and Optimase by Solvay

- 5 Enzymes. Protease enzymes may be incorporated into the compositions in accordance with the invention at a level of from 0.00001% to 2% of enzyme protein by weight of the composition, preferably at a level of from 0.0001% to 1% of enzyme protein by weight of the composition, more preferably at a level of from  
10 0.001% to 0.5% of enzyme protein by weight of the composition, even more preferably at a level of from 0.01% to 0.2% of enzyme protein by weight of the composition.

Lipases: Any lipase suitable for use in alkaline solutions can be used. Suitable lipases include those of bacterial or  
15 fungal origin. Chemically or genetically modified mutants are included.

Examples of useful lipases include a Humicola lanuginosa lipase, e.g., as described in EP 258 068 and EP 305 216, a Rhizomucor miehei lipase, e.g., as described in EP 238 023, a  
20 Candida lipase, such as a C. antarctica lipase, e.g., the C. antarctica lipase A or B described in EP 214 761, a Pseudomonas lipase such as a P. alcaligenes and P. pseudoalcaligenes lipase, e.g., as described in EP 218 272, a P. cepacia lipase, e.g., as described in EP 331 376, a P. stutzeri lipase, e.g., as  
25 disclosed in GB 1,372,034, a P. fluorescens lipase, a Bacillus lipase, e.g., a B. subtilis lipase (Dartois et al., (1993), Biochemica et Biophysica acta 1131, 253-260), a B. stearo-thermophilus lipase (JP 64/744992) and a B. pumilus lipase (WO 91/16422).

30 Furthermore, a number of cloned lipases may be useful, including the Penicillium camembertii lipase described by Yamaguchi et al., (1991), Gene 103, 61-67), the Geotricum candidum lipase (Schimada, Y. et al., (1989), J. Biochem., 106, 383-388), and various Rhizopus lipases such as a R. delemar  
35 lipase (Hass, M.J et al., (1991), Gene 109, 117-113), a R. niveus lipase (Kugimiya et al., (1992), Biosci. Biotech. Biochem. 56, 716-719) and a R. oryzae lipase.

Other types of lipolytic enzymes such as cutinases may also be useful, e.g., a cutinase derived from Pseudomonas

mendocina as described in WO 88/09367, or a cutinase derived from Fusarium solani pisi (e.g. described in WO 90/09446).

Especially suitable lipases are lipases such as M1 Lipase<sup>TM</sup>, Luma fast<sup>TM</sup> and Lipomax<sup>TM</sup> (Genencor), Lipolase<sup>TM</sup> and  
5 Lipolase Ultra<sup>TM</sup> (Novo Nordisk A/S), and Lipase P "Amano" (Amano Pharmaceutical Co. Ltd.).

The lipases are normally incorporated in the detergent composition at a level of from 0.00001% to 2% of enzyme protein by weight of the composition, preferably at a level of from  
10 0.0001% to 1% of enzyme protein by weight of the composition, more preferably at a level of from 0.001% to 0.5% of enzyme protein by weight of the composition, even more preferably at a level of from 0.01% to 0.2% of enzyme protein by weight of the composition.

15 Amylases: Any amylase (a and/or b) suitable for use in alkaline solutions can be used. Suitable amylases include those of bacterial or fungal origin. Chemically or genetically modified mutants are included. Amylases include, for example,  $\alpha$ -amylases obtained from a special strain of B. licheniformis,  
20 described in more detail in GB 1,296,839. Commercially available amylases are Duramyl<sup>TM</sup>, Termamyl<sup>TM</sup>, Fungamyl<sup>TM</sup> and BAN<sup>TM</sup> (available from Novo Nordisk A/S) and Rapidase<sup>TM</sup> and Maxamyl P<sup>TM</sup> (available from Genencor).

The amylases are normally incorporated in the detergent  
25 composition at a level of from 0.00001% to 2% of enzyme protein by weight of the composition, preferably at a level of from 0.0001% to 1% of enzyme protein by weight of the composition, more preferably at a level of from 0.001% to 0.5% of enzyme protein by weight of the composition, even more preferably at a  
30 level of from 0.01% to 0.2% of enzyme protein by weight of the composition.

Cellulases: Any cellulase suitable for use in alkaline solutions can be used. Suitable cellulases include those of bacterial or fungal origin. Chemically or genetically modified  
35 mutants are included. Suitable cellulases are disclosed in US 4,435,307 which discloses fungal cellulases produced from Humicola insolens, in WO 96/34108 and WO 96/34092 which disclose bacterial alkalophilic cellulases (BCE 103) from Bacillus, and in WO 94/21801, US 5,475,101 and US 5,419,778 which disclose EG

III cellulases from *Trichoderma*. Especially suitable cellulases are the cellulases having colour care benefits. Examples of such cellulases are cellulases described in European patent application No. 0 495 257. Commercially available cellulases include Celluzyme<sup>TM</sup> and Carezyme<sup>TM</sup> produced by a strain of *Humicola insolens* (Novo Nordisk A/S), KAC-500(B)<sup>TM</sup> (Kao Corporation), and Puradax<sup>TM</sup> (Genencor International).

Cellulases are normally incorporated in the detergent composition at a level of from 0.00001% to 2% of enzyme protein by weight of the composition, preferably at a level of from 0.0001% to 1% of enzyme protein by weight of the composition, more preferably at a level of from 0.001% to 0.5% of enzyme protein by weight of the composition, even more preferably at a level of from 0.01% to 0.2% of enzyme protein by weight of the composition.

Peroxidases/Oxidases : Peroxidase enzymes are used in combination with hydrogen peroxide or a source thereof (e.g. a percarbonate, perborate or persulfate). Oxidase enzymes are used in combination with oxygen. Both types of enzymes are used for "solution bleaching", i.e. to prevent transfer of a textile dye from a dyed fabric to another fabric when said fabrics are washed together in a wash liquor, preferably together with an enhancing agent as described in e.g. WO 94/12621 and WO 95/01426. Suitable peroxidases/oxidases include those of plant, bacterial or fungal origin. Chemically or genetically modified mutants are included.

Peroxidase and/or oxidase enzymes are normally incorporated in the detergent composition at a level of from 0.00001% to 2% of enzyme protein by weight of the composition, preferably at a level of from 0.0001% to 1% of enzyme protein by weight of the composition, more preferably at a level of from 0.001% to 0.5% of enzyme protein by weight of the composition, even more preferably at a level of from 0.01% to 0.2% of enzyme protein by weight of the composition.

Mixtures of the above mentioned enzymes are encompassed herein, in particular a mixture of a protease, an amylase, a lipase and/or a cellulase.

The enzyme of the invention, or any other enzyme incorporated in the detergent composition, is normally

incorporated in the detergent composition at a level from 0.00001% to 2% of enzyme protein by weight of the composition, preferably at a level from 0.0001% to 1% of enzyme protein by weight of the composition, more preferably at a level from 5 0.001% to 0.5% of enzyme protein by weight of the composition, even more preferably at a level from 0.01% to 0.2% of enzyme protein by weight of the composition.

#### Bleaching agents

10 Additional optional detergent ingredients that can be included in the detergent compositions of the present invention include bleaching agents such as PB1, PB4 and percarbonate with a particle size of 400-800 microns. These bleaching agent components can include one or more oxygen bleaching agents and, 15 depending upon the bleaching agent chosen, one or more bleach activators. When present oxygen bleaching compounds will typically be present at levels of from about 1% to about 25%. In general, bleaching compounds are optional added components in non-liquid formulations, e.g. granular detergents.

20 The bleaching agent component for use herein can be any of the bleaching agents useful for detergent compositions including oxygen bleaches as well as others known in the art.

The bleaching agent suitable for the present invention can be an activated or non-activated bleaching agent.

25 One category of oxygen bleaching agent that can be used encompasses percarboxylic acid bleaching agents and salts thereof. Suitable examples of this class of agents include magnesium monoperoxyphthalate hexahydrate, the magnesium salt of meta-chloro perbenzoic acid, 4-nonylamino-4-oxoperoxybutyric 30 acid and diperoxydodecanedioic acid. Such bleaching agents are disclosed in US 4,483,781, US 740,446, EP 0 133 354 and US 4,412,934. Highly preferred bleaching agents also include 6-nonylamino-6-oxoperoxypropionic acid as described in US 4,634,551.

Another category of bleaching agents that can be used 35 encompasses the halogen bleaching agents. Examples of hypohalite bleaching agents, for example, include trichloro isocyanuric acid and the sodium and potassium dichloroisocyanurates and N-chloro and N-bromo alkane sulphonamides. Such materials are normally added at 0.5-10% by weight of the finished product,

preferably 1-5% by weight.

The hydrogen peroxide releasing agents can be used in combination with bleach activators such as tetra-acetythylenediamine (TAED), nonanoyloxybenzenesulfonate (NOBS, 5 described in US 4,412,934), 3,5-trimethyl-hexsanolxybenzenesulfonate (ISONOBS, described in EP 120 591) or pentaacetylglucose (PAG), which are perhydrolyzed to form a peracid as the active bleaching species, leading to improved bleaching effect. In addition, very suitable are the bleach 10 activators C8(6-octanamido-caproyl) oxybenzene-sulfonate, C9(6-nonanamido caproyl) oxybenzenesulfonate and C10 (6-decanamido caproyl) oxybenzenesulfonate or mixtures thereof. Also suitable activators are acylated citrate esters such as disclosed in European Patent Application No. 91870207.7.

15 Useful bleaching agents, including peroxyacids and bleaching systems comprising bleach activators and peroxygen bleaching compounds for use in cleaning compositions according to the invention are described in application USSN 08/136,626.

The hydrogen peroxide may also be present by adding an 20 enzymatic system (i.e. an enzyme and a substrate therefore) which is capable of generation of hydrogen peroxide at the beginning or during the washing and/or rinsing process. Such enzymatic systems are disclosed in European Patent Application EP 0 537 381.

25 Bleaching agents other than oxygen bleaching agents are also known in the art and can be utilized herein. One type of non-oxygen bleaching agent of particular interest includes photoactivated bleaching agents such as the sulfonated zinc and/or aluminium phthalocyanines. These materials can be deposited 30 upon the substrate during the washing process. Upon irradiation with light, in the presence of oxygen, such as by hanging clothes out to dry in the daylight, the sulfonated zinc phthalocyanine is activated and, consequently, the substrate is bleached. Preferred zinc phthalocyanine and a photoactivated 35 bleaching process are described in US 4,033,718. Typically, detergent composition will contain about 0.025% to about 1.25%, by weight, of sulfonated zinc phthalocyanine.

Bleaching agents may also comprise a manganese catalyst. The manganese catalyst may, e.g., be one of the compounds

described in "Efficient manganese catalysts for low-temperature bleaching", Nature 369, 1994, pp. 637-639.

#### Suds suppressors

5 Another optional ingredient is a suds suppressor, exemplified by silicones, and silica-silicone mixtures. Silicones can generally be represented by alkylated polysiloxane materials, while silica is normally used in finely divided forms exemplified by silica aerogels and xerogels and hydrophobic  
10 silicas of various types. These materials can be incorporated as particulates, in which the suds suppressor is advantageously releasably incorporated in a water-soluble or waterdispersible, substantially non surface-active detergent impermeable carrier. Alternatively the suds suppressor can be dissolved or dispersed  
15 in a liquid carrier and applied by spraying on to one or more of the other components.

A preferred silicone suds controlling agent is disclosed in US 3,933,672. Other particularly useful suds suppressors are the self-emulsifying silicone suds suppressors, described in  
20 German Patent Application DTOS 2,646,126. An example of such a compound is DC-544, commercially available from Dow Corning, which is a siloxane-glycol copolymer. Especially preferred suds controlling agent are the suds suppressor system comprising a mixture of silicone oils and 2-alkyl-alkanols. Suitable 2-alkyl-  
25 alkanols are 2-butyl-octanol which are commercially available under the trade name Isofol 12 R.

Such suds suppressor system are described in European Patent Application EP 0 593 841.

Especially preferred silicone suds controlling agents are  
30 described in European Patent Application No. 92201649.8. Said compositions can comprise a silicone/ silica mixture in combination with fumed nonporous silica such as Aerosil<sup>R</sup>.

The suds suppressors described above are normally employed at levels of from 0.001% to 2% by weight of the composition,  
35 preferably from 0.01% to 1% by weight.

#### Other components

Other components used in detergent compositions may be employed such as soil-suspending agents, soil-releasing agents,

optical brighteners, abrasives, bactericides, tarnish inhibitors, coloring agents, and/or encapsulated or nonencapsulated perfumes.

Especially suitable encapsulating materials are water soluble capsules which consist of a matrix of polysaccharide and polyhydroxy compounds such as described in GB 1,464,616.

Other suitable water soluble encapsulating materials comprise dextrans derived from ungelatinized starch acid esters of substituted dicarboxylic acids such as described in US 3,455,838. These acid-ester dextrans are, preferably, prepared from such starches as waxy maize, waxy sorghum, sago, tapioca and potato. Suitable examples of said encapsulation materials include N-Lok manufactured by National Starch. The N-Lok encapsulating material consists of a modified maize starch and glucose. The starch is modified by adding monofunctional substituted groups such as octenyl succinic acid anhydride.

Antiredeposition and soil suspension agents suitable herein include cellulose derivatives such as methylcellulose, carboxymethylcellulose and hydroxyethylcellulose, and homo- or co-polymeric polycarboxylic acids or their salts. Polymers of this type include the polyacrylates and maleic anhydride-acrylic acid copolymers previously mentioned as builders, as well as copolymers of maleic anhydride with ethylene, methylvinyl ether or methacrylic acid, the maleic anhydride constituting at least 20 mole percent of the copolymer. These materials are normally used at levels of from 0.5% to 10% by weight, more preferably from 0.75% to 8%, most preferably from 1% to 6% by weight of the composition.


Preferred optical brighteners are anionic in character, examples of which are disodium 4,4'-bis-(2-diethanolamino-4-anilino -s- triazin-6-ylamino)stilbene-2:2' disulphonate, disodium 4, - 4'-bis-(2-morpholino-4-anilino-s-triazin-6-ylamino-stilbene-2:2' - disulphonate, disodium 4,4' - bis-(2,4-dianilino-s-triazin-6-ylamino)stilbene-2:2' - disulphonate, monosodium 4',4'' - bis-(2,4-dianilino-s-tri-azin-6-ylamino)stilbene-2-sulphonate, disodium 4,4' -bis-(2-anilino-4-(N-methyl-N-2-hydroxyethylamino)-s-triazin-6-ylamino)stilbene-2,2' - disulphonate, di-sodium 4,4' -bis-(4-phenyl-2,1,3-triazol-2-yl)-stilbene-2,2' disulphonate, di-so-dium 4,4'bis(2-



anilino-4-(1-methyl-2-hydroxyethylamino)-s-triazin-6-ylamino)stilbene-2,2'-disulphonate, sodium 2(stilbyl-4''-(naphtho-1',2':4,5)-1,2,3, - triazole-2''-sulphonate and 4,4'-bis(2-sulphostyryl)biphenyl.

5 Other useful polymeric materials are the polyethylene glycols, particularly those of molecular weight 1000-10000, more particularly 2000 to 8000 and most preferably about 4000. These are used at levels of from 0.20% to 5% more preferably from 0.25% to 2.5% by weight. These polymers and the previously  
10 mentioned homo- or co-polymeric poly-carboxylate salts are valuable for improving whiteness maintenance, fabric ash deposition, and cleaning performance on clay, proteinaceous and oxidizable soils in the presence of transition metal impurities.

Soil release agents useful in compositions of the present  
15 invention are conventionally copolymers or terpolymers of terephthalic acid with ethylene glycol and/or propylene glycol units in various arrangements. Examples of such polymers are disclosed in US 4,116,885 and 4,711,730 and EP 0 272 033. A particular preferred polymer in accordance with EP 0 272 033 has  
20 the formula:



25 where PEG is  $-(\text{OC}_2\text{H}_4)_n-$ , PO is  $(\text{OC}_3\text{H}_6\text{O})$  and T is  $(\text{pOOC}_6\text{H}_4\text{CO})$ .

Also very useful are modified polyesters as random copolymers of dimethyl terephthalate, dimethyl sulfoisophthalate, ethylene glycol and 1,2-propanediol, the end groups consisting primarily of sulphobenzoate and secondarily of  
30 mono esters of ethylene glycol and/or 1,2-propanediol. The target is to obtain a polymer capped at both end by sulphobenzoate groups, "primarily", in the present context most of said copolymers herein will be endcapped by sulphobenzoate groups. However, some copolymers will be less than fully capped,  
35 and therefore their end groups may consist of monoester of ethylene glycol and/or 1,2-propanediol, thereof consist "secondarily" of such species.

The selected polyesters herein contain about 46% by weight of dimethyl terephthalic acid, about 16% by weight of 1,2-

propanediol, about 10% by weight ethylene glycol, about 13% by weight of dimethyl sulfobenzoic acid and about 15% by weight of sulfoisophthalic acid, and have a molecular weight of about 3.000. The polyesters and their method of preparation are described in detail in EP 311 342.

#### Softening agents

Fabric softening agents can also be incorporated into laundry detergent compositions in accordance with the present invention. These agents may be inorganic or organic in type. Inorganic softening agents are exemplified by the smectite clays disclosed in GB-A-1 400898 and in US 5,019,292. Organic fabric softening agents include the water insoluble tertiary amines as disclosed in GB-A1 514 276 and EP 0 011 340 and their combination with mono C<sub>12</sub>-C<sub>14</sub> quaternary ammonium salts are disclosed in EP-B-0 026 528 and di-long-chain amides as disclosed in EP 0 242 919. Other useful organic ingredients of fabric softening systems include high molecular weight polyethylene oxide materials as disclosed in EP 0 299 575 and 0 313 146.

Levels of smectite clay are normally in the range from 5% to 15%, more preferably from 8% to 12% by weight, with the material being added as a dry mixed component to the remainder of the formulation. Organic fabric softening agents such as the water-insoluble tertiary amines or dilong chain amide materials are incorporated at levels of from 0.5% to 5% by weight, normally from 1% to 3% by weight whilst the high molecular weight polyethylene oxide materials and the water soluble cationic materials are added at levels of from 0.1% to 2%, normally from 0.15% to 1.5% by weight. These materials are normally added to the spray dried portion of the composition, although in some instances it may be more convenient to add them as a dry mixed particulate, or spray them as molten liquid on to other solid components of the composition.

35

#### Polymeric dye-transfer inhibiting agents

The detergent compositions according to the present invention may also comprise from 0.001% to 10%, preferably from 0.01% to 2%, more preferably from 0.05% to 1% by weight of

polymeric dye- transfer inhibiting agents. Said polymeric dye-transfer inhibiting agents are normally incorporated into detergent compositions in order to inhibit the transfer of dyes from colored fabrics onto fabrics washed therewith. These  
5 polymers have the ability of complexing or adsorbing the fugitive dyes washed out of dyed fabrics before the dyes have the opportunity to become attached to other articles in the wash.

Especially suitable polymeric dye-transfer inhibiting  
10 agents are polyamine N-oxide polymers, copolymers of N-vinylpyrrolidone and N-vinylimidazole, polyvinylpyrrolidone polymers, polyvinylloxazolidones and polyvinylimidazoles or mixtures thereof.

Addition of such polymers also enhances the performance of  
15 the enzymes according to the invention.

The detergent composition according to the invention can be in liquid, paste, gels, bars or granular forms.

Non-dusting granulates may be produced, e.g., as disclosed in US 4,106,991 and 4,661,452 (both to Novo Industri A/S) and  
20 may optionally be coated by methods known in the art. Examples of waxy coating materials are poly(ethylene oxide) products (polyethyleneglycol, PEG) with mean molecular weights of 1000 to 20000; ethoxylated nonylphenols having from 16 to 50 ethylene oxide units; ethoxylated fatty alcohols in which the alcohol  
25 contains from 12 to 20 carbon atoms and in which there are 15 to 80 ethylene oxide units; fatty alcohols; fatty acids; and mono- and di- and triglycerides of fatty acids. Examples of film-forming coating materials suitable for application by fluid bed techniques are given in GB 1483591.

30 Granular compositions according to the present invention can also be in "compact form", i.e. they may have a relatively higher density than conventional granular detergents, i.e. from 550 to 950 g/l; in such case, the granular detergent compositions according to the present invention will contain a  
35 lower amount of "Inorganic filler salt", compared to conventional granular detergents; typical filler salts are alkaline earth metal salts of sulphates and chlorides, typically sodium sulphate; "Compact" detergent typically comprise not more than 10% filler salt. The liquid compositions according to the

present invention can also be in "concentrated form", in such case, the liquid detergent compositions according to the present invention will contain a lower amount of water, compared to conventional liquid detergents. Typically, the water content of the concentrated liquid detergent is less than 30%, more preferably less than 20%, most preferably less than 10% by weight of the detergent compositions.

The compositions of the invention may for example, be formulated as hand and machine laundry detergent compositions including laundry additive compositions and compositions suitable for use in the pretreatment of stained fabrics, rinse added fabric softener compositions, and compositions for use in general household hard surface cleaning operations and dishwashing operations.

The following examples are meant to exemplify compositions for the present invention, but are not necessarily meant to limit or otherwise define the scope of the invention. In the detergent compositions, the abbreviated component identifications have the following meanings:

20

LAS: Sodium linear C<sub>12</sub> alkyl benzene sulphonate  
TAS: Sodium tallow alkyl sulphate  
XYAS: Sodium C<sub>1X</sub> - C<sub>1Y</sub> alkyl sulfate  
SS: Secondary soap surfactant of formula 2-butyl

25 octanoic acid

25EY: A C<sub>12</sub> - C<sub>15</sub> predominantly linear primary alcohol condensed with an average of Y moles of ethylene oxide

45EY: A C<sub>14</sub> - C<sub>15</sub> predominantly linear primary alcohol condensed with an average of Y moles of ethylene oxide

30 XYEZS: C<sub>1X</sub> - C<sub>1Y</sub> sodium alkyl sulfate condensed with an average of Z moles of ethylene oxide per mole

Nonionic: C<sub>13</sub> - C<sub>15</sub> mixed ethoxylated/propoxylated fatty alcohol with an average degree of ethoxylation of 3.8 and an average degree of propoxylation of 4.5 sold under the tradename

35 Plurafax LF404 by BASF GmbH

CFAA: C<sub>12</sub> - C<sub>14</sub> alkyl N-methyl glucamide

TFAA: C<sub>16</sub> - C<sub>18</sub> alkyl N-methyl glucamide

Silicate: Amorphous Sodium Silicate (SiO<sub>2</sub>:Na<sub>2</sub>O ratio = 2.0)

NaSKS-6: Crystalline layered silicate of formula d-Na<sub>2</sub>Si<sub>2</sub>O<sub>5</sub>

- Carbonate: Anhydrous sodium carbonate  
Phosphate: Sodium tripolyphosphate  
MA/AA: Copolymer of 1:4 maleic/acrylic acid, average molecular weight about 80,000
- 5 Polyacrylate: Polyacrylate homopolymer with an average molecular weight of 8,000 sold under the tradename PA30 by BASF GmbH
- Zeolite A: Hydrated Sodium Aluminosilicate of formula  $\text{Na}_{12}(\text{AlO}_2\text{SiO}_2)_{12} \cdot 27\text{H}_2\text{O}$  having a primary particle size in the
- 10 range from 1 to 10 micrometers
- Citrate: Tri-sodium citrate dihydrate  
Citric: Citric Acid  
Perborate: Anhydrous sodium perborate monohydrate bleach, empirical formula  $\text{NaBO}_2 \cdot \text{H}_2\text{O}_2$
- 15 PB4: Anhydrous sodium perborate tetrahydrate  
Percarbonate: Anhydrous sodium percarbonate bleach of empirical formula  $2\text{Na}_2\text{CO}_3 \cdot 3\text{H}_2\text{O}_2$   
TAED: Tetraacetyl ethylene diamine  
CMC: Sodium carboxymethyl cellulose
- 20 DETPMP: Diethylene triamine penta (methylene phosphonic acid), marketed by Monsanto under the Tradename Dequest 2060  
PVP: Polyvinylpyrrolidone polymer  
EDDS: Ethylenediamine-N, N'-disuccinic acid, [S,S] isomer in the form of the sodium salt
- 25 Suds Suppressor: 25% paraffin wax Mpt 50°C, 17% hydrophobic silica, 58% paraffin oil  
Granular Suds suppressor: 12% Silicone/silica, 18% stearyl alcohol, 70% starch in granular form  
Sulphate: Anhydrous sodium sulphate
- 30 HMWPEO: High molecular weight polyethylene oxide  
TAE 25: Tallow alcohol ethoxylate (25)

#### Detergent Example I

- 35 A granular fabric cleaning composition in accordance with the invention may be prepared as follows:

Sodium linear C <sub>12</sub> alkyl	6.5
benzene sulfonate	

44

	Sodium sulfate	15.0
	Zeolite A	26.0
	Sodium nitrilotriacetate	5.0
	Enzyme of the invention	0.1
5	PVP	0.5
	TAED	3.0
	Boric acid	4.0
	Perborate	18.0
	Phenol sulphonate	0.1
10	Minors	Up to 100

Detergent Example II

A compact granular fabric cleaning composition (density 800 g/l) in accord with the invention may be prepared as

15 follows:

	45AS	8.0
	25E3S	2.0
	25E5	3.0
	25E3	3.0
20	TFAA	2.5
	Zeolite A	17.0
	NaSKS-6	12.0
	Citric acid	3.0
	Carbonate	7.0
25	MA/AA	5.0
	CMC	0.4
	Enzyme of the invention	0.1
	TAED	6.0
	Percarbonate	22.0
30	EDDS	0.3
	Granular suds suppressor	3.5
	water/minors	Up to 100%

Detergent Example III

35 Granular fabric cleaning compositions in accordance with the invention which are especially useful in the laundering of coloured fabrics were prepared as follows:

	LAS	10.7	-
	TAS	2.4	-

45

	TFAA	-	4.0
	45AS	3.1	10.0
	45E7	4.0	-
	25E3S	-	3.0
5	68E11	1.8	-
	25E5	-	8.0
	Citrate	15.0	7.0
	Carbonate	-	10
	Citric acid	2.5	3.0
10	Zeolite A	32.1	25.0
	Na-SKS-6	-	9.0
	MA/AA	5.0	5.0
	DETPMP	0.2	0.8
	Enzyme of the invention	0.10	0.05
15	Silicate	2.5	-
	Sulphate	5.2	3.0
	PVP	0.5	-
	Poly (4-vinylpyridine)-N-Oxide/copolymer of vinyl-	-	0.2
20	imidazole and vinyl-pyrrolidone		
	Perborate	1.0	-
	Phenol sulfonate	0.2	-
	Water/Minors	Up to 100%	

25

Detergent Example IV

Granular fabric cleaning compositions in accordance with the invention which provide "Softening through the wash" capability may be prepared as follows:

30	45AS	-	10.0
	LAS	7.6	-
	68AS	1.3	-
	45E7	4.0	-
	25E3	-	5.0
35	Coco-alkyl-dimethyl hydroxy-ethyl ammonium chloride	1.4	1.0
	Citrate	5.0	3.0
	Na-SKS-6	-	11.0
	Zeolite A	15.0	15.0

46

	MA/AA	4.0	4.0
	DETPMP	0.4	0.4
	Perborate	15.0	-
	Percarbonate	-	15.0
5	TAED	5.0	5.0
	Smectite clay	10.0	10.0
	HMWPEO	-	0.1
	Enzyme of the invention	0.10	0.05
	Silicate	3.0	5.0
10	Carbonate	10.0	10.0
	Granular suds suppressor	1.0	4.0
	CMC	0.2	0.1
	Water/Minors	Up to 100%	

15 Detergent Example V

Heavy duty liquid fabric cleaning compositions in accordance with the invention may be prepared as follows:

	I	II
20 LAS acid form	-	25.0
Citric acid	5.0	2.0
25AS acid form	8.0	-
25AE2S acid form	3.0	-
25AE7	8.0	-
25 CFAA	5	-
DETPMP	1.0	1.0
Fatty acid	8	-
Oleic acid	-	1.0
Ethanol	4.0	6.0
30 Propanediol	2.0	6.0
Enzyme of the invention	0.10	0.05
Coco-alkyl dimethyl hydroxy ethyl ammonium chloride	-	3.0
35 Smectite clay	-	5.0
PVP	2.0	-
Water / Minors	Up to 100%	

Use in the textile and cellulosic fiber processing industries



In the present context, the term "cellulosic material" is intended to mean fibers, sewn and unsewn fabrics, including knits, wovens, denims, yarns, and toweling, made from cotton, cotton blends or natural or manmade cellulosics (e.g. originating from xylan-containing cellulose fibers such as from wood pulp) or blends thereof. Examples of blends are blends of cotton or rayon/viscose with one or more companion material such as wool, synthetic fibers (e.g. polyamide fibers, acrylic fibers, polyester fibers, polyvinyl alcohol fibers, polyvinyl chloride fibers, polyvinylidene chloride fibers, polyurethane fibers, polyurea fibers, aramid fibers), and cellulose-containing fibers (e.g. rayon/viscose, ramie, hemp, flax/linen, jute, cellulose acetate fibers, lyocell).

The preparation of the present invention is useful in the cellulosic fiber processing industry for the pretreatment or retting of fibers from hemp, flax or linen.

The processing of cellulosic material for the textile industry, as for example cotton fiber, into a material ready for garment manufacture involves several steps: spinning of the fiber into a yarn; construction of woven or knit fabric from the yarn and subsequent preparation, dyeing and finishing operations. Woven goods are constructed by weaving a filling yarn between a series of warp yarns; the yarns could be two different types. Knitted goods are constructed by forming a network of interlocking loops from one continuous length of yarn. The cellulosic fibers can also be used for non-woven fabric.

The preparation process prepares the textile for the proper response in dyeing operations. The sub-steps involved in preparation are desizing (for woven goods), scouring and bleaching. A one step combined scour/bleach process is also used by the industry. Although preparation processes are most commonly employed in the fabric state; scouring, bleaching and dyeing operations can also be done at the fiber or yarn stage.

The processing regime can be either batch or continuous with the fabric being contacted by the liquid processing stream in open width or rope form. Continuous operations generally use a saturator whereby an approximate equal weight of chemical bath per weight of fabric is applied to the fabric, followed by a

heated dwell chamber where the chemical reaction takes place. A washing section then prepares the fabric for the next processing step. Batch processing generally takes place in one processing bath whereby the fabric is contacted with approximately 8 -15 times its weight in chemical bath. After a reaction period, the chemicals are drained, fabric rinsed and the next chemical is applied. Discontinuous pad-batch processing involves a saturator whereby an approximate equal weight of chemical bath per weight of fabric is applied to the fabric, followed by a dwell period which in the case of cold pad-batch might be one or more days.

Woven goods are the prevalent form of textile fabric construction. The weaving process demands a "sizing" of the warp yarn to protect it from abrasion. Starch, polyvinyl alcohol (PVA), carboxymethyl cellulose, waxes and acrylic binders are examples of typical sizing chemicals used because of availability and cost. The size must be removed after the weaving process as the first step in preparing the woven goods. The sized fabric in either rope or open width form is brought in contact with the processing liquid containing the desizing agents. The desizing agent employed depends upon the type of size to be removed. For PVA sizes, hot water or oxidative processes are often used. The most common sizing agent for cotton fabric is based upon starch. Therefore most often, woven cotton fabrics are desized by a combination of hot water, the enzyme  $\alpha$ -amylase to hydrolyze the starch and a wetting agent or surfactant. The cellulosic material is allowed to stand with the desizing chemicals for a "holding period" sufficiently long to accomplish the desizing. The holding period is dependent upon the type of processing regime and the temperature and can vary from 15 minutes to 2 hours, or in some cases, several days. Typically, the desizing chemicals are applied in a saturator bath which generally ranges from about 15°C to about 55°C. The fabric is then held in equipment such as a "J-box" which provides sufficient heat, usually between about 55°C and about 100°C, to enhance the activity of the desizing agents. The chemicals, including the removed sizing agents, are washed away from the fabric after the termination of the holding period.

In order to ensure a high whiteness or a good wettability

and resulting dyeability, the size chemicals and other applied chemicals must be thoroughly removed. It is generally believed that an efficient desizing is of crucial importance to the following preparation processes: scouring and bleaching.

5       The scouring process removes much of the non-cellulosic compounds naturally found in cotton. In addition to the natural non-cellulosic impurities, scouring can remove dirt, soils and residual manufacturing introduced materials such as spinning, coning or slashing lubricants. The scouring process employs  
10 sodium hydroxide or related causticizing agents such as sodium carbonate, potassium hydroxide or mixtures thereof. Generally an alkali stable surfactant is added to the process to enhance solubilization of hydrophobic compounds and/or prevent their redeposition back on the fabric. The treatment is generally at a  
15 high temperature, 80°C - 100°C, employing strongly alkaline solutions, pH 13-14, of the scouring agent. Due to the non-specific nature of chemical processes not only are the impurities but the cellulose itself is attacked, leading to damages in strength or other desirable fabric properties. The  
20 softness of the cellulosic fabric is a function of residual natural cotton waxes. The non-specific nature of the high temperature strongly alkaline scouring process cannot discriminate between the desirable natural cotton lubricants and the manufacturing introduced lubricants. Furthermore, the  
25 conventional scouring process can cause environmental problems due to the highly alkaline effluent from these processes. The scouring stage prepares the fabric for the optimal response in bleaching. An inadequately scoured fabric will need a higher level of bleach chemical in the subsequent bleaching stages.

30       The bleaching step decolorizes the natural cotton pigments and removes any residual natural woody cotton trash components not completely removed during ginning, carding or scouring. The main process in use today is an alkaline hydrogen peroxide bleach. In many cases, especially when a very high whiteness is  
35 not needed, bleaching can be combined with scouring.

It is contemplated that the scouring step can be carried out using the xyloglucanase or xyloglucanase preparation of the present invention in combination with a few other enzyme activities at a temperature of about 50°C - 80°C and a pH of

about 7-11, thus substituting or supplementing the highly causticizing agents.

#### MATERIALS AND METHODS

##### 5 Strains:

*Bacillus licheniformis*, ATCC 14580, and *Bacillus agaradhaerens*, NCIMB 40482, respectively, comprises a DNA sequence of the invention encoding a xyloglucanase.

##### 10 Other strains:

*E. coli* strain: Cells of *E. coli* SJ2 (Diderichsen et al., 1990) were prepared for and transformed by electroporation using a Gene Pulser<sup>TM</sup> electroporator from BIO-RAD as described by the supplier.

15 *B. subtilis* PL1885. (Diderichsen et al., (1990)).

*B. subtilis* PL2306. This strain is the *B. subtilis* DN1885 with disrupted *apr* and *npr* genes (Diderichsen et al. (1990)) disrupted in the transcriptional unit of the known *Bacillus subtilis* cellulase gene, resulting in cellulase negative cells.

20 *B. subtilis* PL2316. This strain is the *B. subtilis* DN1885 with disrupted *apr* and *npr* genes (Diderichsen, B., Wedsted, U., Hedegaard, L., Jensen, B. R., Sjøholm, C. (1990) Cloning of *aldB*, which encodes alpha-acetolactate decarboxylase, an exoenzyme from *Bacillus brevis*. J. Bacteriol., 172, 4315-4321)  
25 disrupted in the transcriptional unit of the known *Bacillus subtilis* xylanase gene, resulting in xylanase negative cells. The disruptions were performed essentially as described in A.L. Sonenshein et al. (1993).

Competent cells were prepared and transformed as described by  
30 Yasbin et al. (1975).

#### Plasmids.

pSJ1678 (see WO 94/19454).

pmOL944. This plasmid is a pUB110 derivative essentially  
35 containing elements making the plasmid propagatable in *Bacillus subtilis*, kanamycin resistance gene and having a strong promoter and signal peptide cloned from the *amyL* gene of *B. licheniformis* ATCC 14580. The signal peptide contains a SacII site making it

convenient to clone the DNA encoding the mature part of a protein in-fusion with the signal peptide. This results in the expression of a Pre-protein which is directed towards the exterior of the cell.

- 5       The plasmid was constructed by means of ordinary genetic engineering and is briefly described in the following.

Construction of pMOL944:

The pUB110 plasmid (McKenzie, T. et al., 1986,) was digested with the unique restriction enzyme NciI. A PCR fragment  
10 amplified from the amyL promoter encoded on the plasmid pDN1981 (Jørgensen et al., 1990) was digested with NciI and inserted in the NciI digested pUB110 to give the plasmid pSJ2624.

The two PCR primers used have the following sequences:

# LWN5494 5'-GTCGCCGGGGCGGCCGCTATCAATTGGTAACTGTATCTCAGC -3'  
15 # LWN5495 5'-GTCGCCGGGGAGCTCTGATCAGGTACCAAGCTTGTCGACCTGCAGAA  
TGAGGCAGCAAGAAGAT -3'

The primer #LWN5494 inserts a NotI site in the plasmid.

The plasmid pSJ2624 was then digested with SacI and NotI and a new PCR fragment amplified on amyL promoter encoded on the  
20 pDN1981 was digested with SacI and NotI and this DNA fragment was inserted in the SacI-NotI digested pSJ2624 to give the plasmid pSJ2670.

This cloning replaces the first amyL promoter cloning with the same promoter but in the opposite direction. The two primers  
25 used for PCR amplification have the following sequences:

#LWN5938 5'-GTCGGCGGGCCGCTGATCACGTACCAAGCTTGTCGACCTGCAGAAATG  
AGGCAGCAAGAAGAT -3'  
#LWN5939 5'-GTCGGAGCTCTATCAATTGGTAACTGTATCTCAGC -3'

The plasmid pSJ2670 was digested with the restriction  
30 enzymes PstI and BclI and a PCR fragment amplified from a cloned DNA sequence encoding the alkaline amylase SP722 (International Patent Application published as WO95/26397 which is hereby incorporated by reference) was digested with PstI and BclI and inserted to give the plasmid pMOL944. The two primers used for  
35 PCR amplification have the following sequence:

#LWN7864 5' -AACAGCTGATCAGCACTGATCTTTTAGCTTGGCAC-3'  
#LWN7901 5' -AACTGCAGCCGCGGCACATCATAATGGGACAAATGGG -3'

The primer #LWN7901 inserts a SacII site in the plasmid.

**Media:**

TY (as described in Ausubel, F. M. et al. 1995).

LB agar (as described in Ausubel, F. M. et al, 1995).

5 LBPG is LB agar supplemented with 0.5% Glucose and 0.05 M potassium phosphate, pH 7.0

AZCL-xyloglucan is added to LBPG-agar to 0.5 % .AZCL-xyloglucan is from Megazyme, Australia.

BPX media is described in EP 0 506 780 (WO 91/09129).

10 **Medium A:** Per flask: 30 g wheat bran, 45 ml of the following solution: 10 g rofec (Roquette 101-0441), 10 g  $\text{NH}_4\text{NO}_3$  (Merck 1187), 10 g  $\text{KH}_2\text{PO}_4$  (Merck 4873), 40 g Solcafloc (Dicacel available from Dicalite-Europe-Nord, 9000 Gent, Belgium), 0.75 g  $\text{MgSO}_4 \cdot 7\text{H}_2\text{O}$  (Merck 5886), 15 g  $\text{CaCO}_3$ , tap water to 1000 ml, pH  
15 adjusted to 6.5. Autoclave for 40 min at 121° C.

**Medium B:** 30 g soyabean meal, 15 g maltodex 01 (Roquette 101-7845), 5 g peptone (Difco 0118), 0.2 ml pluronic (PE-6100, 101-3068), deionized water up to 1000 ml. 100 ml in 500 ml Erlenmeyer flask with 2 baffles. Autoclave at 121° C for 40 min.

20 **Medium C:** 15 g wheat bran, 5 g dextrose, 6.7 g Bacto Yeast Nitrogen Base, deionized water up to 1000 ml. 100 ml in 500 ml Erlenmeyer flask with 2 baffles. Autoclave at 121° C for 40 min.

**Medium D:** 20% sucrose, 5% soy flakes and 1% sodium phosphate.

25 **Medium E:** 5 g Yeast Extract, 10 g Tryptone, 3 g  $(\text{NH}_4)_2\text{SO}_4$ , 3 g  $\text{K}_2\text{HPO}_4$ , 2 g  $\text{KH}_2\text{PO}_4$ , 1 g CMC, 10 g maltodextrin, 30 g wheat bran, deionized water up to 1000 ml, pH adjusted to 7.0. 100 ml in 500 ml Erlenmeyer flask with 2 baffles. Autoclave at 121° C for 40 min.

30

**Fermentation procedure:**

The fungal strains were grown in shake flasks under the following growth conditions:

35 **Media:** A, B or C (see list of media)

**Temperature:** 26°C

**RPM:** A, stationary

B and C, 125 - 200

**Incubation time:** A, 6 - 30 days

B and C, 2 - 21 days

Bacteria were grown in shake flasks containing medium D or E at 30°C with 250 rpm shaking for 3-4 days.

5

**Xyloglucanase assay (XGU):**

The xyloglucanase activity is measured using AZCL-xyloglucan from Megazyme, Australia, as substrate.

A solution of 0.2 % of the blue substrate is suspended in a 10 0.1 M phosphate buffer pH 7.5 under stirring. The solution is distributed under stirring to 1.5 ml Eppendorf tubes (0.75 ml to each), 50 µl enzyme solution is added and they are incubated in an Eppendorp Thermomixer model 5436 for 20 min. at 40°C with a mixing of 1200 rpm. After incubation the coloured solution is 15 separated from the solid by 4 min. centrifugation at 14,000 rpm and the absorbance of the supernatant is measured at 600 nm.

One XGU units is defined as the amount of enzyme resulting in an absorbance of 0.24 in a 1 cm cuvette at 600 nm.

20 **Isoelectric focusing:**

Isoelectric focusing was carried out in precast Ampholine PAG plates pH 3.5-9.5 (Pharmacia, Sweden) according to the manufacturer's instructions. The samples were applied in duplicate and after electrophoresis the gel was divided into 25 two. An overlay containing 1% agarose and 0.4% AZCL xyloglucan in water was poured onto one half of the gel, a similar overlay containing AZCL HE cellulose was poured onto the other half. Incubation at 30°C for 2-16 hours. Enzyme activity was identified by blue zones.

30

**General molecular biology methods:**

DNA manipulations and transformations were performed using standard methods of molecular biology (Sambrook et al. (1989) Molecular cloning: A laboratory manual, Cold Spring Harbor lab., 35 Cold Spring Harbor, NY; Ausubel, F. M. et al. (eds.) "Current protocols in Molecular Biology". John Wiley and Sons, 1995; Harwood, C. R., and Cutting, S. M. (eds.) "Molecular Biological Methods for Bacillus". John Wiley and Sons, 1990).

Enzymes for DNA manipulations were used according to the

specifications of the suppliers.

The following examples illustrate the invention.

5 EXAMPLE 1

**Screening for microorganisms producing alkaline xyloglucanase**

The microorganisms to be screened were grown in a liquid culture as described in the Materials and Methods section. After centrifugation the culture supernatants were tested for  
10 xyloglucanase and cellulase activity. The following assays were used: agarose plates containing 1% agarose in 0.08 M Britton-Robinson buffer pH 7 or pH 9, and 0.2% AZCL xyloglucan and 0.2% AZCL HE cellulose, respectively, 10 ml samples were applied into  
15 16 holes in the agarose plates, incubation at 30°C for 2-16 hours. Enzyme activity was identified by blue halos.

Culture broths with good activity on AZCL xyloglucan and no or low activity on AZCL HE cellulose were further tested by isoelectric focusing (IEF) as described in the Materials and Methods section. The microorganisms listed in the Materials and  
20 Methods section were found to produce an enzyme with activity on AZCL xyloglucan which by IEF was separable from activity on AZCL HE cellulose. When screening for xyloglucanase activity also cellulase activity was found, and it was determined whether both activities derives from the same enzyme. This was done by  
25 separation of the enzymes by IEF isoelectric focusing which separates the protein according to the charge or pI. After this separation the different enzymatic activities was again identified using overlay techniques. The sample was run at two parallel wells: one stained for xyloglucanase and another for  
30 AZCL HE- cellulose. If the same lane has both activities it is a cellulase, if only one lane has xyloglucanase and no cellulase activity it is a xyloglucanase which can be further characterized either by purification or by cloning.

35 EXAMPLE 2

**Production of a *Bacillus licheniformis* xyloglucanase (XG)**

This example illustrates a method for producing xyloglucanase from *Bacillus licheniformis*.

*Bacillus licheniformis*, ATCC 14580, was grown in shake



flasks using the substrate PS 1 (20% sucrose and 5% soy flakes and 1% sodium phosphate) at 30°C with 250 rpm shaking for 4 days. The culture broth (total 10 litres) was adjusted to pH 7.5 with NaOH, followed by treatment with 50 ml of a cationic  
5 flocculation agent under stirring at room temperature and subsequently 470 ml of a 0.1% solution of an anionic flocculation agent. The flocculated material was separated by centrifugation using a Sorval RC 3B 10.000 rpm for 30 min. The supernatant was clarified using a Whatman glass filter number F.  
10 A total of 9 litres was obtained with an activity of 10 XGU per ml.

The liquid was concentrated to 1.5 litres on a filtron with a cut off at 10 kDa.

Bacitracin cross linked to Sepharose was used for affinity  
15 column chromatography for removal of the proteases. Then the not bound material was adjusted to pH 5.0 using acetic acid and applied to a SP-Sepharose column adjusted with 20 mM sodium acetate buffer pH 5.0. The XG activity was eluted from the column by using a 0.5 M NaCl gradient. The fractions containing  
20 XG activity was pooled and concentrated on an Amicon cell with a GR 81 polysulfon membrane with a cut off at 8 kDa.

The concentrated solution containing 363 XGU per ml was applied to size chromatography using a Superdex 200 column equilibrated with a buffer of 0.1 M sodium acetate pH 6.0. The  
25 pure XG 1 eluted with an apparent molecular weight of 26 kDa and gave a single band in SDS-PAGE of 26 kDa. The specific activity was determined to 221 XGU per A.280.

### EXAMPLE 3

#### 30 Characterisation of a *Bacillus licheniformis* xyloglucanase

The amino acid sequence of the enzyme produced as described in Example 2 was obtained after SDS-PAGE and electroblotting of the 26 kDa protein. The amino acid sequence is listed in the appended SEQ ID No:2.

35 This amino acid shows highest homology with glycosyl hydrolases from Family 12 which at present are classified as endo-beta-1,4-glucanases (EC 3.2.1.4): 65% homology with the family 12 *Erwinia carotovora*  $\beta$ -1,4-glucan glucanohydrolase (cels - P16630 Swissprot).

Accordingly, the present invention further relates to an enzyme which has the amino acid sequence listed in SEQ ID NO:2 or has an amino acid sequence which is at least 70%, preferably at least 75%, more preferably at least 80%, more preferably at least 85%, even more preferably at least 90%, especially at least 95%, homologous therewith.

The obtained *Bacillus licheniformis* xyloglucanase enzyme was analysed using Megazyme AZCL xyloglucan for xyloglucanase activity determination and AZCL HE cellulose for endoglucanase activity determination. The relative release of blue colour was 40 times higher on the xyloglucan substrate compared with the HE cellulose using the same amount of purified enzyme.

The same result was found for the acidic *Aspergillus aculeatus* xyloglucanase enzyme (EG II) described in the patent application WO 94/14953 which also showed 2% relative activity against HE cellulose.

#### Temperature optimum

The AZCL xyloglucan from Megazyme was also used for determination of temperature optimum. The activity was measured at different temperatures after incubation with the xyloglucanase of Example 2 for 20 minutes and the release of blue colour was determined at 600 nm. The major part of the colour was released at 60°C, but still 50% relative activity was obtained at 70° C.

#### pH activity profile

For obtaining a realistic pH activity profile, the activity determinations were carried out using buffers having a pKa value within 1.0 of the actual pH. The following buffer systems were used: pH 4-5.5 sodium acetate, pH 6 Mes buffer (Sigma), pH 6.5 - 7.5 Mops buffer (Sigma), pH 8-8.5 Barbiturate, pH 9-10.5 Glycine.

The pH was measured after incubation in a parallel handled tube. Incubation 20 min. at 40° C. Final substrate concentration 1.33 gram xyloglucan per l. This is the  $K_M$  (see below for steady state at pH 7.5). The activity was determined after measurement of the formation of reducing ends as described for steady state kinetic.

pH	% relative activity
4.13	7
4.59	32
5.05	83
5.54	100
6.00	97
6.49	84
6.99	96
7.52	90
7.99	61
8.49	45
9.02	27
9.41	14
9.90	5

**EXAMPLE 4****5 Comparison example: Steady state kinetics on soluble xyloglucan and CMC**

A method for determination of activity against xyloglucan has been developed.

The substrate is xyloglucan (amyloid) from tamarind seeds  
10 (the substrate is commercially available from Megazyme). Buffer 0.1 M sodium phosphate pH 7.5.

The substrate is prepared as a stock solution containing 5 gram per l in buffer. After mixing it is heated using a magnetic stirrer until a clear solution is obtained. The solution is then  
15 cooled to 40 °C and kept in a temperature controlled water bath at 40 °C.

The diluted enzyme solution of 0.5 ml is preheated for 10 min. and mixed with 1.0 ml substrate and incubated for 20 min.

The formation of reducing sugars is determined by using p-  
20 hydroxy-benzoic-acid-hydrazide (PHBAH) modified from Lever (1972) using 5 gram of potassium sodium tartrate in addition to 1.5 gram of PHBAH. Glucose is used as reference for

determination of the reducing groups.

The apparent catalytic properties of 2 known cellulases (endo-beta-1,4-glucanases) on xyloglucan from tamarind seeds was measured:

5 a. EG I from *Humicola insolens* (classified as belonging to family 7 of glycosyl hydrolases) disclosed in WO 95/24471.

b. EG III from *Trichoderma* (classified as belonging to family 12 of glycosyl hydrolases) disclosed in US patent US 5,475,101.

10 All the enzymes were purified to high homogeneity giving a single band in SDS-PAGE and the molar extinction coefficient was used for calculation of the enzyme concentration. The determination was based on different concentrations of the substrate from 0.25 to 3.3 gram per litre. The kinetic  
15 determination was according to the computer program Grafit and assuming Michelis Menten kinetic.

### Results

Enzyme of the invention: *Bacillus licheniformis*

20 Xyloglucanase 1 with a molecular weight (MW) of 26 kDa. Based on a molar absorbance of 78,000, determined by amino acid analysis, the  $k_{cat}$  was 16.5 per sec (std. error 0.6) on xyloglucan at pH 7.5,  $K_m$  1.1 g/l (std. error 0.1). Using CMC as substrate the  $k_{cat}$  was impossible to measure but below 3 per sec. The ratio of  
25 maximum xyloglucanase activity to maximum activity on CMC is at least 5:1.

a. Comparison EG I, alkaline cellulase from *Humicola insolens*, MW 50 kDa, molar extinction coefficient of 66310. A  $k_{cat}$  of 19 per sec (std. error 0.7) on xyloglucan at pH 7.5,  $K_m$   
30 0.7 g/l (std. error 0.08). On CMC the  $k_{cat}$  is 86 per sec (std. error 5). The ratio of maximum xyloglucanase activity to maximum activity on CMC is 2:9.

b. Comparison EG III, acid cellulase from *Trichoderma*, MW 24 kDa, molar extinction coefficient of 71930. A  $k_{cat}$  of 16 per  
35 sec (std. error 1.7) on xyloglucan at pH 7.5,  $K_m$  0.5 g/l (std. error 0.15). On CMC the  $k_{cat}$  is 18 per sec (Std. error 0.6). The ratio of maximum xyloglucanase activity to maximum activity on CMC is 8:9.

## EXAMPLE 5

**Cloning and expression of the *Bacillus licheniformis* xyloglucanase gene**5 Genomic DNA preparation:

Strain *Bacillus licheniformis*, ATCC 14580, was propagated in liquid TY medium. After 16 hours incubation at 30°C and 300 rpm, the cells were harvested, and genomic DNA isolated by the method described by Pitcher et al., (1989).

10 Genomic library construction:

Genomic DNA was partially digested with restriction enzyme Sau3A, and size-fractionated by electrophoresis on a 0.7 % agarose gel. Fragments between 2 and 10 kb in size was isolated by electrophoresis onto DEAE-cellulose paper (Dretzen et al.,  
15 (1981)).

Isolated DNA fragments were ligated to BamHI digested pSJ1678 plasmid DNA, and the ligation mixture was used to transform *E. coli* SJ2.

Identification of positive clones:

20 A DNA library in *E. coli*, constructed as described above, was screened on LB agar plates containing 0.5% AZCL-xyloglucan (Megazyme) and 9 µg/ml Chloramphenicol and incubated overnight at 37°C. Clones expressing hydrolysing activity appeared with blue diffusion halos. Positive clones were plated on LB agar  
25 plates containing 0.5% AZCL-xyloglucan (Megazyme). The plasmids of this clones were isolated by Qiagen plasmid spin preps on 1 ml of overnight culture broth (cells incubated at 37°C in TY with 9 µg/ml Chloramphenicol and shaking at 250 rpm). One of these clones (PL2949) was further characterized by DNA  
30 sequencing of the cloned Sau3A DNA fragment.

The DNA was characterised by DNA sequencing using the Taq deoxy-terminal cycle sequencing kit (Perkin-Elmer, USA), fluorescent labelled terminators and appropriate oligonucleotides as primers.

35 Analysis of the sequence data was performed according to Devereux et al. (1984). The sequence encoding the mature protein is shown in SEQ ID NO:1 (signal plus mature; the mature part corresponding to positions 88-783. The derived protein sequence

is shown in SEQ ID NO:2, the mature protein corresponding to positions 30-261 of SEQ ID NO:2.

Subcloning and expression of the xyloglucanase gene from  
5 *B.licheniformis* in *B.subtilis*:

The xyloglucanase encoding DNA sequence of the invention was PCR amplified using the PCR primer set consisting of these two oligonucleotides:

10 Xyloglu .upper.PstI

5'-GCCTCATTCTGCAGCAGCGGCGGCTTCGTCATCAAACCCGTCGG-3'

Xyloglu .lower.NotI

5'-GCTGCATCGGCATCGCGGCCGCGCAATACGTAAGGATGGTATCG -3'

15

Restriction sites PstI and NotII are underlined.

Chromosomal DNA isolated from *B. licheniformis* as described above was used as template in a PCR reaction using Amplitaq DNA Polymerase (Perkin Elmer) according to  
20 manufacturers instructions. The PCR reaction was set up in PCR buffer (10 mM Tris-HCl, pH 8.3, 50 mM KCl, 1.5 mM MgCl<sub>2</sub>, 0.01 % (w/v) gelatin) containing 200 µM of each dNTP, 2.5 units of AmpliTaQ polymerase (Perkin-Elmer, Cetus, USA) and 100 pmol of each primer  
25 The PCR reactions was performed using a DNA thermal cycler (Landgraf, Germany). One incubation at 94°C for 1 min followed by thirty cycles of PCR performed using a cycle profile of denaturation at 94°C for 30 sec, annealing at 60°C for 1 min, and extension at 72 °C for 2 min. Five-µl aliquots of the ampli-  
30 fication product was analysed by electrophoresis in 0.7 % agarose gels (NuSieve, FMC). The appearance of a DNA fragment size approx. 0.8 kb indicated proper amplification of the gene segment.

35 Subcloning of PCR fragment:

Fortyfive-µl aliquots of the PCR products generated as described above were purified using QIAquick PCR purification kit (Qiagen, USA) according to the manufacturer's instructions. The purified DNA was eluted in 50 µl of 10mM Tris-HCl, pH 8.5.

5  $\mu$ g of pMOL944 and twentyfive- $\mu$ l of the purified PCR fragment was digested with PstI and NotI, electrophoresed in 0.8 % low gelling temperature agarose (SeaPlaque GTG, FMC) gels, the relevant fragments were excised from the gels, and purified  
5 using QIAquick Gel extraction Kit (Qiagen, USA) according to the manufacturer's instructions. The isolated PCR DNA fragment was then ligated to the PstI-NotI digested and purified pMOL944. The ligation was performed overnight at 16°C using 0.5  $\mu$ g of each DNA fragment, 1 U of T4 DNA ligase and T4 ligase buffer  
10 (Boehringer Mannheim, Germany).

The ligation mixture was used to transform competent *B. subtilis* PL2316. The transformed cells were plated onto LBPG- 10  $\mu$ g/ml of Kanamycin-agar plates. After 18 hours incubation at 37°C colonies were seen on plates. Several clones were analyzed  
15 by isolating plasmid DNA from overnight culture broth.

One such positive clone was restreaked several times on agar plates as used above, this clone was called PL2954. The clone PL2954 was grown overnight in TY-10 $\mu$ g/ml Kanamycin at 37°C, and next day 1 ml of cells were used to isolate plasmid  
20 from the cells using the Qiaprep Spin Plasmid Miniprep Kit #27106 according to the manufacturers recommendations for *B. subtilis* plasmid preparations. This DNA was DNA sequenced and revealed the DNA sequence corresponding to the mature part of the xyloglucanase in the SEQ ID NO. 1 which is shown in  
25 positions 88-783.

Expression and purification of *B. licheniformis* yloglucanase:

PL2954 was grown in 25 x 200 ml BPX media with 10  $\mu$ g/ml of Kanamycin in 500 ml two baffled shakeflasks for 5 days at 37°C  
30 at 300 rpm.

**EXAMPLE 6**

**Cloning and expression of the *Bacillus agaradhaerens* xyloglucanase gene**

35 Genomic DNA preparation:

Strain *Bacillus agaradhaerens*, NCIMB 40482, was propagated in liquid medium as described in WO94/01532. After 16 hours incubation at 30°C and 300 rpm, the cells were harvested, and genomic DNA isolated by the method described by Pitcher et al.

(1989).

Genomic library construction:

Genomic DNA was partially digested with restriction enzyme  
Sau3A, and size-fractionated by electrophoresis on a 0.7 %  
5 agarose gel. Fragments between 2 and 7 kb in size was isolated  
by electrophoresis onto DEAE-cellulose paper (Dretzen, G.,  
Bellard, M., Sassone-Corsi, P., Chambon, P. (1981) A reliable  
method for the recovery of DNA fragments from agarose and  
acrylamide gels. Anal. Biochem., 112, 295-298).

10 Isolated DNA fragments were ligated to BamHI digested  
pSJ1678 plasmid DNA, and the ligation mixture was used to  
transform *E. coli* SJ2.

Cells were plated on LB agar plates containing 0.1% CMC  
(Sodium-Carboxy-Methyl-Cellulose, Aqualon, France) and 9 µg/ml  
15 Chloramphenicol and incubated overnight at 37°C.

Identification of positive clones:

A DNA library in *E. coli*, constructed as described above,  
was screened on LB agar plates containing 0.1% CMC (Sodium-  
Carboxy-Methyl-Cellulose, Aqualon, France) and 9 µg/ml  
20 Chloramphenicol and incubated overnight at 37°C. The  
transformants were subsequently replica plated onto the same  
type of plates, and these new plates were incubated 8 hours or  
overnight at 37°C.

The original plates were coloured using 1 mg/ml of Congo  
25 Red (SIGMA, USA). The coloring was continued for half an hour  
with moderate orbital shaking, after which the plates were  
washed two times 15 minutes using 1 M NaCl.

Yellowish halos appeared at positions where cellulase  
positive clones were present, from the replica plates these  
30 cellulase positive clones were rescued and restreaked onto LB  
agar plates containing 0.1% CMC and 9 µg/ml Chloramphenicol and  
incubated overnight at 37°C.

One such clone (MB110) was further characterized by DNA  
sequencing of the cloned Sau3A DNA fragment.

35 The DNA was characterised by DNA sequencing by  
primerwalking, using the Taq deoxy-terminal cycle sequencing kit  
(Perkin-Elmer, USA), fluorescent labelled terminators and  
appropriate oligonucleotides as primers.



Analysis of the sequence data was performed according to Devereux et al. (1984). The sequence encoding the mature protein, subcloned in the example below, is shown in SEQ ID NO:1. The derived protein sequence is shown in SEQ ID NO:2.

5 Subcloning and expression of xyloglucanase in *B.subtilis*:

The xyloglucanase encoding DNA sequence of the invention was PCR amplified using the PCR primer set consisting of these two oligo nucleotides:

10 Xyloglucanase.upper.SacII

5'-CAT TCT GCA GCC GCG GCA GAA GAT GTC ACT TCG TCA CAG -3'

Xyloglucanase.lower.NotI

5'-GTT GAG AAA AGC GGC CGC CAC TTC TAA AGT TCT AAA GCA CG -3'

15 Restriction sites SacII and NotI are underlined.

Chromosomal DNA isolated from *B.agaradherans* as described above was used as template in a PCR reaction using Amplitaq DNA Polymerase (Perkin Elmer) according to manufacturers instructions. The PCR reaction was set up in PCR  
20 buffer (10 mM Tris-HCl, pH 8.3, 50 mM KCl, 1.5 mM MgCl<sub>2</sub>, 0.01 % (w/v) gelatin) containing 200 µM of each dNTP, 2.5 units of AmpliTaq polymerase (Perkin-Elmer, Cetus, USA) and 100 pmol of each primer

The PCR reactions was performed using a DNA thermal  
25 cycler (Landgraf, Germany). One incubation at 94°C for 1 min followed by thirty cycles of PCR performed using a cycle profile of denaturation at 94°C for 30 sec, annealing at 60°C for 1 min, and extension at 72 °C for 2 min. Five-µl aliquots of the amplification product was analysed by electrophoresis in 0.7 %  
30 agarose gels (NuSieve, FMC). The appearance of a DNA fragment size 1.6 kb indicated proper amplification of the gene segment.  
Subcloning of PCR fragment:

Fortyfive-µl aliquots of the PCR products generated as described above were purified using QIAquick PCR purification  
35 kit (Qiagen, USA) according to the manufacturer's instructions. The purified DNA was eluted in 50 µl of 10mM Tris-HCl, pH 8.5. 5 µg of pMOL944 and twentyfive-µl of the purified PCR fragment was digested with SacII and NotI, electrophoresed in 0.8 % low gelling temperature agarose (SeaPlaque GTG, FMC) gels, the

relevant fragments were excised from the gels, and purified using QIAquick Gel extraction Kit (Qiagen, USA) according to the manufacturer's instructions. The isolated PCR DNA fragment was then ligated to the SacII-NotI digested and purified pMOL944.  
5 The ligation was performed overnight at 16°C using 0.5 µg of each DNA fragment, 1 U of T4 DNA ligase and T4 ligase buffer (Boehringer Mannheim, Germany).

The ligation mixture was used to transform competent *B. subtilis* PL2306. The transformed cells were plated onto LBPG-  
10 10 µg/ml of Kanamycin -0.1% AZCL-Xyloglucan-agar plates. After 18 hours incubation at 37°C cells positively expressing the cloned Xyloglucanase were seen as colonies surrounded by blue halos. One such positive clone was restreaked several times on agar plates as used above, this clone was called MB563. The  
15 clone MB563 was grown overnight in TY-10µg/ml Kanamycin at 37°C, and next day 1 ml of cells were used to isolate plasmid from the cells using the Qiaprep Spin Plasmid Miniprep Kit #27106 according to the manufacturers recommendations for *B. subtilis* plasmid preparations.

20 Expression and purification of *B. agaradhaerens* xyloglucanase:

MB563 was grown in 25 x 200 ml BPX media with 10 µg/ml of Kanamycin in 500 ml two baffled shakeflasks for 5 days at 37°C at 300 rpm.

25 **EXAMPLE 7**

**Characterisation of a *Bacillus agaradhaerens* xyloglucanase**

Purification and characterization:

7000 ml shake flask culture fluid from *Bacillus* with the  
30 clone MB 563 expressed as described in example 6 was received. The fermentation medium was adjusted to pH 7.5 with NaOH and flocculated using cationic flocculation agent C521 (10% solution) and 0.1% solution of anionic agent A130: To 7000 ml of fermentation medium was added 168 ml of C521 (10%) simultaneous  
35 with 335 ml of A130 under stirring at room temperature. The flocculated material was separated by centrifugation using a Sorval RC 3B centrifuge at 10,000 rpm for 30 minutes. The supernatant was clarified using Whatman glass filter number F. In total was obtained 6500 ml of clear solution containing

227,500 XGU units.

The clear solution of 2500 ml was applied to a 1000 ml Q-Sepharose column equilibrated with 50 mM Tris buffer pH 7.0. The bound enzyme was eluted using a NaCl gradient.

5 The partly purified product was concentrated using an Amicon ultrafiltration cell with a membrane with a cut-off value of 6 kDa. Total 70.000 XGU units and 2.5 gram of enzyme protein was obtained.

The pure sample gave a single band in SDS-PAGE with a  
10 apparent molecular weight of 61 kDa. A molar extinction coefficient of 123,040 was used for calculation of enzyme protein concentration and is based on the amino acid composition deducted from the DNA sequence.

The concentrated fraction was formulated with 40% sorbitol  
15 and used for enzyme trials in detergent and textile applications.

#### Immunological methods:

Highly purified *Bacillus agaradhaerens* XEG1 obtained from  
20 clone MB 563 was used for production of antiserum.

The immunization procedure was conducted at DAKO using rabbits. Each rabbit was immunized with 100  $\mu$ l cellulase (0.4 mg protein per ml) mixed with 100  $\mu$ l adjuvant. Each rabbit was immunized 15 times with one weeks interval. The rabbit serum was  
25 collected and the gammaglobulin purified from the serum.

#### Temperature optimum:

The AZCL xyloglucan from Megazyme was also used for determination of temperature optimum. The activity was measured  
30 at different temperatures after incubation with the *Bacillus agaradhaerens* XEG1 for 20 minutes and the release of blue colour was determined at 600 nm. The major part of the colour was released at 50°C, but still 20% relative activity was obtained at 60°C.

35

#### Steady state kinetics on soluble xyloglucan and CMC:

The method for determination of activity against xyloglucan described in example 4 was applied to the *B. agaradhaerens*

xyloglucanase of the invention. The substrate was diluted to at least 8 different concentrations with 4 below the apparent  $K_m$  (see below).

The following apparent catalytic properties of the *Bacillus* 5 *agaradhaerens* XEG1 (endo-beta-1,4-xylo-glucanases) on xyloglucan from tamarind seeds were found: At pH 7.5 the  $k_{cat}$  of 183 per sec and a apparent  $K_m$  of 0.05 gram per l was obtained. For comparison the activity on CMC (Degree of substitution 0.7 and a degree of polymerization of 200) using the same steady state 10 kinetic method with 8 different substrate concentrations in duplicate below the  $K_m$  the following data was obtained:  $k_{cat}$  of 64 per sec and a  $K_m$  of 2.2 gram per l, indicating that this enzyme prefer xyloglucan from carboxymethyl cellulose.

Alkaline activity, at pH 10 using a glycine buffer and 15 xyloglucan substrate, resulted in a  $K_{cat}$  of 90 per sec and a apparent  $K_m$  of 0.08 gram per l indicate that this xyloglucanase has a very high alkaline activity.

The pH activity profile using blue Megazyme AZCL xyloglucan indicate that the enzyme has more than 50% relative activity in 20 the pH interval of 5.0 to 10.5.

In a detergent matrices using the blue Megazyme substrate the following data was obtained. In US Tide powder detergent 1 gram per l with 9 German hardness of water 66% relative activity to buffer pH 7.5. Using European conditions and powder Ariel in 25 5 gram per l and 18 German hardness the relative activity of 86% was obtained. These data indicate that the *Bacillus* *agaradhaerens* XEG1 is well suited to be used in detergent matrices.

## LITERATURE

- Ausubel, F. M. et al. (eds.) "Current protocols in Molecular Biology". John Wiley and Sons, 1995.
- 5 N. C. Carpita and D. M. Gibeaut (1993) *The Plant Journal* 3:1-30.
- T. Christensen et al. *Biotechnology* vol 6 page 1419-1422, 1988.
- 10 Devereux et al. (1984) *Nucleic Acids Res.* 12, 387-395.
- Diderichsen, B., Wedsted, U., Hedegaard, L., Jensen, B. R., Sjøholm, C. (1990) Cloning of aldB, which encodes alpha-acetolactate decarboxylase, an exoenzyme from *Bacillus brevis*.
- 15 *J. Bacteriol.* 172:4315-4321.
- Dretzen, G., Bellard, M., Sassone-Corsi, P., Chambon, P. (1981) A reliable method for the recovery of DNA fragments from agarose and acrylamide gels. *Anal. Biochem.*, 112, 295-298.
- 20 Eriksson, O.E. & Hawksworth, D.L.: *Systema Ascomycetum* vol 12 (1993).
- S. C. Fry et al (1992) *Biochemical Journal* 282:821-828
- 25 Hawksworth, D.L., Kirk, P.M., Sutton, B.C. and Pegler, D.N.: *Dictionary of the fungi*, International Mycological Institute, 616 pp (1995);
- 30 T. Hayashi and D. P. Delmer (1988) *Carbohydrate Research* 181:273-277.
- Henrissat, B. 1991. A classification of glycosyl hydrolases based on amino acid sequence similaritites. *Biochem. J.*,
- 35 280:309-316.
- Henrissat, B., and A. Bairoch. 1993. New families in the classification of glycosyl hydrolases based on amino acid sequence similaritites. *Biochem. J.*, 293:781-788.

- Jülich, W.: Higher Taxa of Basidiomycetes, *Bibliotheca Mycologia* 85, 485 pp (1981).
- 5 Jørgensen, P.L. et al., 1990, *Gene*, 96, p. 37-41.
- McKenzie, T. et al., 1986, *Plasmid* 15:93-103.
- Leatherbarrow, R. J. (1992) *Grafit* version 3.0 Erithacus  
10 Software Ltd. Staines, U.K.
- Lever, M. (1972) A new reaction for colormetric determination of carbohydrates. *Anal. Biochem.* 47, 273-279.
- 15 O'Donnell, K.: *Zygomycetes in culture*, University of Georgia, US, 257 pp (1979).
- Pitcher, D. G., Saunders, N. A., Owen, R. J. (1989). Rapid extraction of bacterial genomic DNA with guanidium thiocyanate.  
20 *Lett. Appl. Microbiol.*, 8, 151-156.
- J. K. C. Rose et al (1996) *Plant Physiology* 110:493-499
- A.L. Sonenshein, J.A. Hoch and Richard Losick (Eds.) (1993)  
25 *Bacillus subtilis* and other Gram-Positive Bacteria, American Society for microbiology, p.618.
- Vincken, J.P., Beldman, G., and Voragen, A.G.J. Substrate-specificity of endoglucanases - what determines xyloglucanase  
30 activity. *Carbohydrate Research* 298(4):299-310, 1997.
- Von Arx, J. A.: The genera of fungi sporulating in culture, 424 pp (1981).
- 35 R. L. Whistler and J. N. BeMiller (1993) *Industrial gums: Polysaccharides and their derivatives*, Academic Press Inc.
- Yasbin, R.E., Wilson, G.A. and Young, F.E. (1975) Transformation and transfection in lysogenic strains of *Bacillus subtilis* :

evidence for selective induction of prophage in competent cells.  
J. Bacteriol, 121:296-304.

W.S. York et al (1996) Carbohydrate Research 285:99-128.

## SEQUENCE LISTING

## (1) GENERAL INFORMATION:

- 5       (i) APPLICANT:
- (A) NAME: NOVO NORDISK A/S
  - (B) STREET: Novo Alle
  - (C) CITY: Bagsvaerd
  - (E) COUNTRY: Denmark
  - 10       (F) POSTAL CODE (ZIP): DK-2880
  - (G) TELEPHONE: +45 44 44 88 88
  - (H) TELEFAX: +45 44 49 32 56
- 15       (ii) TITLE OF INVENTION: ALKALINE XYLOGLUCANASE
- (iii) NUMBER OF SEQUENCES: 4
- (iv) COMPUTER READABLE FORM:
- (A) MEDIUM TYPE: Floppy disk
  - 20       (B) COMPUTER: IBM PC compatible
  - (C) OPERATING SYSTEM: PC-DOS/MS-DOS
  - (D) SOFTWARE: PatentIn Release #1.0, Version #1.30
- (EPO)
- 25 (2) INFORMATION FOR SEQ ID NO: 1:
- (i) SEQUENCE CHARACTERISTICS:
- (A) LENGTH: 786 base pairs
  - (B) TYPE: nucleic acid
  - 30       (C) STRANDEDNESS: single
  - (D) TOPOLOGY: linear
- (ii) MOLECULE TYPE: DNA (genomic)
- 35       (vi) ORIGINAL SOURCE:
- (A) ORGANISM: Bacillus licheniformis ATCC 14580
- (xi) SEQUENCE DESCRIPTION: SEQ ID NO: 1:



71

GTGAAAAACA ACCATTTGCT AAAATCCATT CTGCTATGGG GTGCCGTATG CATCATAGTG 60  
 CTGGCAGGGC CGCTCTCAGC ATTTGCCGCT TCGTCATCAA ACCCGTCGGA TAAATTGTAT 120  
 5 TTTAAAAACA AAAAATACTA CATATTCAAC AATGTATGGG GAGCCGACCA GGTCAGCGGC 180  
 TGGTGGCAGA CCATTTATCA TAATAGTGAT TCAGATATGG GCTGGGTGTG GAATTGGCCG 240  
 AGCAATACAA GCACGGTAAA AGCTTATCCG TCGATCGTCA GCGGCTGGCA TTGGACTGAA 300  
 10 GGCTATACTG CCGGAAGCGG CTTCCCGACG CGATTGTCAG ATCAAAAAA CATCAACACG 360  
 AAAGTCAGCT ATTCGATCAG CGCAAACGGC ACATACAATG CCGCATATGA CATTGGCTC 420  
 15 CACAATACAA ACAAGGCGAG CTGGGATTCG GCTCCAACCG ATGAGATTAT GATCTGGCTC 480  
 AATAACACAA ACGCCGGACC TGCCGGTTCC TATGTCGAAA CTGTATCGAT TGGCGGGCAC 540  
 AGTTGGAAAG TATATAAAGG CTATATTGAT GCTGGAGGCG GCAAAGGGTG GAACGTGTTT 600  
 20 TCATTTATCA GAACAGCAAA CACCCAAAGT GCGAACCTGA ATATTCGGA TTTCACGAAT 660  
 TATCTTGCCG ACTCCAAACA GTGGCTTTCC AAAACAAAGT ATGTCAGCAG TGTGGAATTC 720  
 25 GGTACTGAAG TTTTCGGAGG CACAGGACAA ATTAATATT CCAATTGGGA CGTAACGGTC 780  
 CGCTGA 786

(2) INFORMATION FOR SEQ ID NO: 2:

30

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 261 amino acids
- (B) TYPE: amino acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

35

(ii) MOLECULE TYPE: protein

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 2:

40

Val Lys Asn Asn His Leu Leu Lys Ser Ile Leu Leu Trp Gly Ala Val  
 1                      5                      10                      15  
 Cys Ile Ile Val Leu Ala Gly Pro Leu Ser Ala Phe Ala Ala Ser Ser

72

	20	25	30
	Ser Asn Pro Ser Asp Lys Leu Tyr Phe Lys Asn Lys Lys Tyr Tyr Ile		
	35	40	45
5	Phe Asn Asn Val Trp Gly Ala Asp Gln Val Ser Gly Trp Trp Gln Thr		
	50	55	60
	Ile Tyr His Asn Ser Asp Ser Asp Met Gly Trp Val Trp Asn Trp Pro		
10	65	70	75 80
	Ser Asn Thr Ser Thr Val Lys Ala Tyr Pro Ser Ile Val Ser Gly Trp		
	85	90	95
15	His Trp Thr Glu Gly Tyr Thr Ala Gly Ser Gly Phe Pro Thr Arg Leu		
	100	105	110
	Ser Asp Gln Lys Asn Ile Asn Thr Lys Val Ser Tyr Ser Ile Ser Ala		
	115	120	125
20	Asn Gly Thr Tyr Asn Ala Ala Tyr Asp Ile Trp Leu His Asn Thr Asn		
	130	135	140
	Lys Ala Ser Trp Asp Ser Ala Pro Thr Asp Glu Ile Met Ile Trp Leu		
25	145	150	155 160
	Asn Asn Thr Asn Ala Gly Pro Ala Gly Ser Tyr Val Glu Thr Val Ser		
	165	170	175
30	Ile Gly Gly His Ser Trp Lys Val Tyr Lys Gly Tyr Ile Asp Ala Gly		
	180	185	190
	Gly Gly Lys Gly Trp Asn Val Phe Ser Phe Ile Arg Thr Ala Asn Thr		
	195	200	205
35	Gln Ser Ala Asn Leu Asn Ile Arg Asp Phe Thr Asn Tyr Leu Ala Asp		
	210	215	220
	Ser Lys Gln Trp Leu Ser Lys Thr Lys Tyr Val Ser Ser Val Glu Phe		
40	225	230	235 240
	Gly Thr Glu Val Phe Gly Gly Thr Gly Gln Ile Asn Ile Ser Asn Trp		
	245	250	255
45	Asp Val Thr Val Arg		
	260		

## (2) INFORMATION FOR SEQ ID NO: 3:

## (i) SEQUENCE CHARACTERISTICS:

- 5 (A) LENGTH: 1614 base pairs  
(B) TYPE: nucleic acid  
(C) STRANDEDNESS: single  
(D) TOPOLOGY: linear

## (ii) MOLECULE TYPE: DNA (genomic)

10

## (vi) ORIGINAL SOURCE:

- (A) ORGANISM: *Bacillus agaradhaerens* NCIMB 40482

## (xi) SEQUENCE DESCRIPTION: SEQ ID NO: 3:

15

GAAGATGTCA CTTGTCACA GTTGGATATT CACTCCTATG TAGCTGACAT GCAGCCTGGC 60  
TGGAATTTAG GAAATACGTT TGACGCTGTT GGAGATGATG AAACAGCGTG GGGGAATCCT 120  
20 CGTGTAAACAA GAGAGTTAAT AAAACGATT GCTGATGAAG GGTATAAAAG CATTGATATC 180  
CCAGTGACAT GGCAAAATCA AATGGGTGGT TCTCCAGATT ATACGATAAA TGAAGATTAT 240  
ATCAATCGGG TGGAGCAAGC GATAGATTGG GCGTTGGAGG AAGACTTATA TGTGATGTTA 300  
25 AATGTGCATC ATGACTCATG GCTGTGGATG TATGATATGG AACATAACTA TGATGAGGTC 360  
ATGGCAAGAT ATACAGCTAT TTGGGAACAA TTGTCGGAAA AATTCAAAG CCACTCCCAT 420  
30 AAGTTGATGT TTGAGAGTGT CAATGAGCCT AGGTTTACGC AGGAGTGGGG AGAGATTCAA 480  
GAAAATCATC ATGCTTACTT AGAAGATTTA AATAAGACGT TCTATTATAT TGTCAGAGAG 540  
TCAGGAGGCA ATAATGTGGA GCGCCCTTTA GTATTGCCTA CGATAGAAAC AGCCACGTCT 600  
35 CAGGATTTAC TAGATCGCTT GTATCAAACA ATGGAAGACT TGGATGATCC TTATTTAATT 660  
GCCACGGTGC ATTATTATGG CTTCTGGCCA TTAGTGTCA ATATAGCAGG GTACACTCAT 720  
40 TTTGAACAGG AAACACAACA AGATATTATA GACACCTTTG ACCGTGTTCA TAACACATTT 780  
ACAGCGCGTG GTGTCCAGT TGTATTAGGC GAATTCGGTT TGTTAGGCTT TGACAAAAGT 840  
ACGGATGTGA TTCAGCAAGG GGAGAAATTA AAGTTTTTTG AGTTTCTCAT CCATCATCTC 900

AATGAACGTG ATATAACCCA TATGTTATGG GATAACGGCC AGCATTTTAA TCGAGAACT 960  
TATGCATGGT ATGATCAAGA ATTCATGAC ATATTAAAAG CGAGTTGGGA GGGGCGTTCT 1020  
5 GCTACAGCAG AGTCTAATTT GATTTCATGTG AAGGACGGAA AGCCAATTAG AGATCAAGAT 1080  
ATACAGCTTT ACTTAAACGG AAATGAGCTA ACAGCCTTAC AGGCAGGGGA GGAATCGCTT 1140  
10 GTTCTAGGAG AGGATTATGA ACTAGCAGGA GCGGTATTAA CGCTAAAAGC GGACACCCTC 1200  
ACAAGACTAA TTACCCCTGG TCAATTAGGA ACCAATGCAG TCATCACAGC ACAATTTAAT 1260  
TCTGGAGCAG ACTGGCGTTT TCAATTACAG AATGTGGACG TGCCAACGGT CGAAAATACA 1320  
15 GATGGCTCAA CATGGCATT TCGATCCCT ACCCATTTTA ATGGTGATAG TCTTGCGACG 1380  
ATGGAAGCTG TTTATGCAA CGGAGAATAT GCTGGGCCGC AAGATTGGAC GTCATTTAAA 1440  
20 GAATTTGGCG AGGCGTTTTC TCCTAATTAC CCCACAGGGG AAATTATTAT ATCAGAAGCC 1500  
TTCTTTAACG CCGTACGGGA TGATGATATC CATTTAACAT TTCATTTTGT GAGCGGAGAG 1560  
ACGGTGGAAT ATACCTTACG TAAAAATGGC AATTATGTTT AAGGTAGACG GTAA 1614  
25

(2) INFORMATION FOR SEQ ID NO: 4:

(i) SEQUENCE CHARACTERISTICS:

- 30 (A) LENGTH: 537 amino acids  
(B) TYPE: amino acid  
(C) STRANDEDNESS: single  
(D) TOPOLOGY: linear

35 (ii) MOLECULE TYPE: protein

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 4:

40 Glu Asp Val Thr Ser Ser Gln Leu Asp Ile His Ser Tyr Val Ala Asp  
1 5 10 15  
Met Gln Pro Gly Trp Asn Leu Gly Asn Thr Phe Asp Ala Val Gly Asp  
20 25 30  
Asp Glu Thr Ala Trp Gly Asn Pro Arg Val Thr Arg Glu Leu Ile Lys

	35	75	40	45	
	Thr Ile Ala Asp Glu Gly Tyr Lys Ser Ile Arg Ile Pro Val Thr Trp				
	50	55	60		
5	Gln Asn Gln Met Gly Gly Ser Pro Asp Tyr Thr Ile Asn Glu Asp Tyr				
	65	70	75	80	
	Ile Asn Arg Val Glu Gln Ala Ile Asp Trp Ala Leu Glu Glu Asp Leu				
10		85	90	95	
	Tyr Val Met Leu Asn Val His His Asp Ser Trp Leu Trp Met Tyr Asp				
	100	105	110		
15	Met Glu His Asn Tyr Asp Glu Val Met Ala Arg Tyr Thr Ala Ile Trp				
	115	120	125		
	Glu Gln Leu Ser Glu Lys Phe Lys Ser His Ser His Lys Leu Met Phe				
20	130	135	140		
	Glu Ser Val Asn Glu Pro Arg Phe Thr Gln Glu Trp Gly Glu Ile Gln				
	145	150	155	160	
	Glu Asn His His Ala Tyr Leu Glu Asp Leu Asn Lys Thr Phe Tyr Tyr				
25		165	170	175	
	Ile Val Arg Glu Ser Gly Gly Asn Asn Val Glu Arg Pro Leu Val Leu				
	180	185	190		
30	Pro Thr Ile Glu Thr Ala Thr Ser Gln Asp Leu Leu Asp Arg Leu Tyr				
	195	200	205		
	Gln Thr Met Glu Asp Leu Asp Asp Pro Tyr Leu Ile Ala Thr Val His				
	210	215	220		
35	Tyr Tyr Gly Phe Trp Pro Phe Ser Val Asn Ile Ala Gly Tyr Thr His				
	225	230	235	240	
	Phe Glu Gln Glu Thr Gln Gln Asp Ile Ile Asp Thr Phe Asp Arg Val				
40		245	250	255	
	His Asn Thr Phe Thr Ala Arg Gly Val Pro Val Val Leu Gly Glu Phe				
	260	265	270		
45	Gly Leu Leu Gly Phe Asp Lys Ser Thr Asp Val Ile Gln Gln Gly Glu				
	275	280	285		

76

Lys Leu Lys Phe Phe Glu Phe Leu Ile His His Leu Asn Glu Arg Asp  
 290 295 300

5 Ile Thr His Met Leu Trp Asp Asn Gly Gln His Phe Asn Arg Glu Thr  
 305 310 315 320

Tyr Ala Trp Tyr Asp Gln Glu Phe His Asp Ile Leu Lys Ala Ser Trp  
 325 330 335

10 Glu Gly Arg Ser Ala Thr Ala Glu Ser Asn Leu Ile His Val Lys Asp  
 340 345 350

Gly Lys Pro Ile Arg Asp Gln Asp Ile Gln Leu Tyr Leu Asn Gly Asn  
 355 360 365

15 Glu Leu Thr Ala Leu Gln Ala Gly Glu Glu Ser Leu Val Leu Gly Glu  
 370 375 380

20 Asp Tyr Glu Leu Ala Gly Gly Val Leu Thr Leu Lys Ala Asp Thr Leu  
 385 390 395 400

Thr Arg Leu Ile Thr Pro Gly Gln Leu Gly Thr Asn Ala Val Ile Thr  
 405 410 415

25 Ala Gln Phe Asn Ser Gly Ala Asp Trp Arg Phe Gln Leu Gln Asn Val  
 420 425 430

Asp Val Pro Thr Val Glu Asn Thr Asp Gly Ser Thr Trp His Phe Ala  
 435 440 445

30 Ile Pro Thr His Phe Asn Gly Asp Ser Leu Ala Thr Met Glu Ala Val  
 450 455 460

35 Tyr Ala Asn Gly Glu Tyr Ala Gly Pro Gln Asp Trp Thr Ser Phe Lys  
 465 470 475 480

Glu Phe Gly Glu Ala Phe Ser Pro Asn Tyr Ala Thr Gly Glu Ile Ile  
 485 490 495

40 Ile Ser Glu Ala Phe Phe Asn Ala Val Arg Asp Asp Asp Ile His Leu  
 500 505 510

Thr Phe His Phe Trp Ser Gly Glu Thr Val Glu Tyr Thr Leu Arg Lys  
 515 520 525

45 Asn Gly Asn Tyr Val Gln Gly Arg Arg  
 530 535

## CLAIMS

1. An enzyme preparation comprising a xyloglucanase having a relative xyloglucanase activity of at least 50% at pH 7.
- 5 2. The preparation according to claim 1, which has a relative xyloglucanase activity of at least 60%, preferably at least 75%, more preferably at least 80%, especially at least 90%, at pH 7.
- 10 3. The preparation according to claim 1 or 2, wherein, at pH 7, the relative xyloglucanase activity is higher than the relative cellulase activity towards carboxymethylcellulose (CMC activity) derived from the xyloglucanase.
- 15 4. The preparation according to any of the claims 1-3, which has a relative xyloglucanase activity of at least 50%, preferably at least 60%, more preferably at least 75%, especially at least 90%, at pH 8.
- 20 5. The preparation according to claim 4, wherein, at pH 8, the relative xyloglucanase activity is higher than the relative cellulase activity towards carboxymethylcellulose (CMC activity) derived from the xyloglucanase.
- 25 6. The preparation according to any of the claims 1-5, wherein the xyloglucanase has a minor or no activity on cellulose or cellulose derivative substrates.
- 30 7. The preparation according to claim 6, wherein the cellulose or cellulose derivative substrate is carboxymethylcellulose (CMC) or microcrystalline cellulose (Avicel).
8. The preparation according to claim 7, wherein the ratio of maximum xyloglucanase activity to maximum activity on CMC or  
35 Avicel is at least 2:1.
9. The preparation according to claim 8, wherein the ration is at least 4:1, preferably at least 5:1, more preferably at least 8:1, especially at least 10:1.

10. The preparation according to any of the claims 1-9, which is derived or derivable from a microorganism, preferably from a bacterium or a fungus, including a yeast.
- 5 11. The preparation according to claim 10, which is derived or derivable from a bacterium, preferably from a Gram positive bacterium, preferably from the *Bacillus/Lactobacillus* subdivision, preferably a strain from the genus *Bacillus*,  
10 especially a strain of *Bacillus licheniformis*, e.g. *Bacillus licheniformis*, ATCC 14580.
12. The preparation according to claim 10 or 11, which is derived or derivable from an alkalophilic bacterium, preferably  
15 from the strain *Bacillus agaradhaerens*, especially *B. agaradhaerens*, NCIMB 40482.
13. The preparation according to any of the claims 1-12, wherein the xyloglucanase belongs to family 7 of glycosyl hydrolases.
- 20 14. The preparation according to any of the claims 1-12, wherein the xyloglucanase belongs to family 12 of glycosyl hydrolases.
15. The preparation according to any of the claims 1-12, wherein  
25 the xyloglucanase belongs to family 5 of glycosyl hydrolases.
16. The preparation according to claim 14 which is derived or derivable by *Bacillus licheniformis*, ATCC 14580.
- 30 17. The preparation according to claim 15 which is derived or derivable by *Bacillus agaradhaerens*, NCIMB 40482.
18. A xyloglucanase which is  
(a) a polypeptide produced by *Bacillus agaradhaerens*, NCIMB  
35 40482, or  
(b) a polypeptide comprising an amino acid sequence as shown in positions 1-537 SEQ ID NO:4, or  
(c) an analogue of the polypeptide defined in (a) or (b) which is at least 70% homologous with said polypeptide, or is derived



from said polypeptide by substitution, deletion or addition of one or several amino acids, or is immunologically reactive with a polyclonal antibody raised against said polypeptide in purified form.

5

19. A xyloglucanase which is

(a) a polypeptide produced by *Bacillus licheniformis*, ATCC 14580, or

(b) a polypeptide comprising an amino acid sequence as shown in positions 30-261 of SEQ ID NO:2, or

(c) an analogue of the polypeptide defined in (a) or (b) which is at least 70% homologous with said polypeptide, or is derived from said polypeptide by substitution, deletion or addition of one or several amino acids, or is immunologically reactive with a polyclonal antibody raised against said polypeptide in purified form.

20. An isolated polynucleotide molecule encoding a polypeptide having xyloglucanase activity selected from the group consisting of:

(a) polynucleotide molecules comprising a nucleotide sequence as shown in SEQ ID NO:1 from nucleotide 88 to nucleotide 783;

(b) polynucleotide molecules that encode a polypeptide that is at least 70% identical to the amino acid sequence of SEQ ID NO:2 from amino acid residue 30 to amino acid residue 261; and

(c) degenerate nucleotide sequences of (a) or (b).

21. An isolated polynucleotide molecule encoding a polypeptide having xyloglucanase activity selected from the group consisting of:

(a) polynucleotide molecules comprising a nucleotide sequence as shown in SEQ ID NO:3 from nucleotide 1 to nucleotide 1611;

(b) polynucleotide molecules that encode a polypeptide that is at least 70% identical to the amino acid sequence of SEQ ID NO:4 from amino acid residue 1 to amino acid residue 537; and

(c) degenerate nucleotide sequences of (a) or (b).

22. The isolated polynucleotide molecule according to claim 20 or 21, wherein the polynucleotide is DNA.

23. An isolated polynucleotide molecule encoding a polypeptide having xyloglucanase activity which polynucleotide molecule hybridizes to a denatured double-stranded DNA probe under medium stringency conditions, wherein the probe is selected from the group consisting of DNA probes comprising the sequence shown in positions 88-783 of SEQ ID NO:1 and DNA probes comprising a subsequence of positions 88-783 of SEQ ID NO:1 having a length of at least about 100 base pairs.

10

24. An isolated polynucleotide molecule encoding a polypeptide having xyloglucanase activity which polynucleotide molecule hybridizes to a denatured double-stranded DNA probe under medium stringency conditions, wherein the probe is selected from the group consisting of DNA probes comprising the sequence shown in positions 1-1611 of SEQ ID NO:3 and DNA probes comprising a subsequence of positions 1-1611 of SEQ ID NO:3 having a length of at least about 100 base pairs.

20 25. An expression vector comprising the following operably linked elements: a transcription promoter; a DNA segment selected from the group consisting of (a) polynucleotide molecules encoding a polypeptide having xyloglucanase activity comprising a nucleotide sequence as shown in SEQ ID NO:1 from nucleotide 88 to nucleotide 783, (b) polynucleotide molecules encoding a polypeptide having xyloglucanase activity that is at least 70% identical to the amino acid sequence of SEQ ID NO:2 from amino acid residue 30 to amino acid residue 261, and (c) degenerate nucleotide sequences of (a) or (b); and a transcription terminator.

26. An expression vector comprising the following operably linked elements: a transcription promoter; a DNA segment selected from the group consisting of (a) polynucleotide molecules encoding a polypeptide having xyloglucanase activity comprising a nucleotide sequence as shown in SEQ ID NO:1 from nucleotide 1 to nucleotide 1611, (b) polynucleotide molecules encoding a polypeptide having xyloglucanase activity that is at least 70% identical to the amino acid sequence of SEQ ID NO:2

from amino acid residue 1 to amino acid residue 537, and (c) degenerate nucleotide sequences of (a) or (b); and a transcription terminator.

5 27. A cultured cell into which has been introduced an expression vector according to claim 25 or 26, wherein said cell expresses the polypeptide encoded by the DNA segment.

28. An isolated polypeptide having xyloglucanase activity  
10 selected from the group consisting of:  
    (a) polypeptide molecules comprising an amino acid sequence as shown in SEQ ID NO: 2 from residue 30 to residue 261; and  
    (b) polypeptide molecules that are at least 70% identical to the amino acids of SEQ ID NO: 2 from amino acid residue 30 to amino  
15 acid residue 261.

29. The polypeptide according to claim 28 which is produced by *Bacillus licheniformis*.

20 30. An isolated polypeptide having xyloglucanase activity selected from the group consisting of:  
    (a) polypeptide molecules comprising an amino acid sequence as shown in SEQ ID NO: 4 from amino acid residue 1 to amino acid residue 537; and  
25 (b) polypeptide molecules that are at least 70% identical to the amino acids of SEQ ID NO: 4 from residue 1 to residue 537.

31. The polypeptide according to claim 30 which is produced by *Bacillus agaradherens*.

30

32. An enzyme preparation comprising a purified polypeptide according to claim 29 or 30.

33. A method of producing a polypeptide having xyloglucanase  
35 activity comprising culturing a cell into which has been introduced an expression vector according to claim 25 or 26, whereby said cell expresses a polypeptide encoded by the DNA segment; and recovering the polypeptide.

34. The preparation according to claim 32 which further comprises one or more enzymes selected from the group consisting of proteases, cellulases (endoglucanases),  $\beta$ -glucanases, hemicellulases, lipases, peroxidases, laccases,  $\alpha$ -amylases, 5 glucoamylases, cutinases, pectinases, reductases, oxidases, phenoloxidases, ligninases, pullulanases, pectate lyases, xyloglucanases, xylanases, pectin acetyl esterases, polygalacturonases, rhamnogalacturonases, pectin lyases, other mannanases, pectin methylesterases, cellobiohydrolases, 10 transglutaminases; or mixtures thereof.

35. An isolated enzyme having xyloglucanase activity, in which the enzyme is (i) free from homologous impurities, and (ii) produced by the method according to claim 33.

15

36. A detergent composition comprising the enzyme preparation according to any of the claims 1-17, 32 or 34 or the enzyme according to any of the claims 18, 19, 28-31 and 35.

20 37. A process for machine treatment of fabrics which process comprises treating fabric during a washing cycle of a machine washing process with a washing solution containing the enzyme preparation according to any of the claims 1-17, 32 or 34 or the enzyme according to any of the claims 18, 19, 28-31 and 35.

25

38. Use of the enzyme preparation according to any of the claims 1-17, 32 or 34 or the enzyme according to any of the claims 18, 19, 28-31 and 35 in the textile industry for improving the properties of cellulosic fibres, yarn, woven or non-woven 30 fabric.

39. The use according to claim 29, wherein the enzyme preparation or the enzyme is used in a scouring process step.

35 40. Use of the enzyme preparation according to any of the claims 1-17, 32 or 34 or the enzyme according to any of the claims 18, 19, 28-31 and 35 in the cellulose fiber processing industry for ratting of fibers selected from the group consisting of hemp, jute, flax and linen.

## INTERNATIONAL SEARCH REPORT

International application No.  
PCT/DK 98/00290

## A. CLASSIFICATION OF SUBJECT MATTER

IPC6: C12N 9/42 // C12N 9/42, C12R 1:07

According to International Patent Classification (IPC) or to both national classification and IPC

## B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

IPC6: C12N

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

SE,DK,FI,NO classes as above

Electronic data base consulted during the international search (name of data base and, where practicable, search terms used)

WPI, CA, BIOSIS, MEDLINE, STRAND

## C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	Patent Abstracts of Japan, abstract of JP 50-68549 A (AJINOMOTO CO INC), 23 March 1993 (23.03.93)	1-3
	--	
X	WO 9634092 A2 (GENENCOR INTERNATIONAL, INC.), 31 October 1996 (31.10.96)	1,2,4,18,24, 36-40
	--	
X	AUST. J. PLANT PHYSIOL., Volume 19, 1992, Gordon MacLachlan et al, "Multiple Forms of 1, 4-beta-Glucanase in Ripening Tomato Fruits I nclude a Xyloglucanase Activatable by Xyloglucan Oligosaccharides", page 137 - page 146, figure 1	1,2,4
	--	

☒ Further documents are listed in the continuation of Box C.☒ See patent family annex.

## \* Special categories of cited documents:

- "A" document defining the general state of the art which is not considered to be of particular relevance
- "E" earlier document but published on or after the international filing date
- "L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)
- "O" document referring to an oral disclosure, use, exhibition or other means
- "P" document published prior to the international filing date but later than the priority date claimed

"T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention

"X" document of particular relevance: the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone

"Y" document of particular relevance: the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art

"&amp;" document member of the same patent family

Date of the actual completion of the international search

26 October 1998

Date of mailing of the international search report

30 October 1998 (30.10.98)

Name and mailing address of the ISA/  
Swedish Patent Office  
Box 5055, S-102 42 STOCKHOLM  
Facsimile No. +46 8 666 02 86

Authorized officer

Hampus Rystedt  
Telephone No. +46 8 782 25 00

## INTERNATIONAL SEARCH REPORT

International application No.

PCT/DK 98/00290

C (Continuation). DOCUMENTS CONSIDERED TO BE RELEVANT		
Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
A	WO 9317101 A1 (UNILEVER PLC ET AL), 2 Sept 1993 (02.09.93)  --	1-40
A	SYMBIOSIS, Volume 21, 1996, A. Rejon-Palomares et al, "Presence of Xyloglucan-Hydrolyzing Glucanases (Xylogucanases) in Arbuscular Mycorrhizal Symbiosis", page 249 - page 261, figure 1, abstract  -- -----	1-17